



REVIEW

Global research priorities for historical ecology to inform conservation

Loren McClenachan^{1,2,*,#}, Torben Rick³, Ruth H. Thurstan⁴, Andrew Trant⁵, Peter S. Alagona⁶, Heidi K. Alleway⁷, Chelsey Armstrong⁸, Rebecca Bliege Bird⁹, Nadia T. Rubio-Cisneros^{10,11}, Miguel Clavero¹², André C. Colonese¹³, Katie Cramer¹⁴, Ancilleno O. Davis¹⁵, Joshua Drew¹⁶, Michelle M. Early-Capistrán¹⁷, Graciela Gil-Romera¹⁸, Molly Grace¹⁹, Marco B. A. Hatch²⁰, Eric Higgs², Kira Hoffman^{21,22}, Jeremy B. C. Jackson²³, Antonieta Jerardino²⁴, Michelle J. LeFebvre²⁵, Heike K. Lotze²⁶, Ryan S. Mohammed²⁷, Naia Morueta-Holme²⁸, Catalina Munteanu²⁹, Alexis M. Mychajliw³⁰, Bonnie Newsom³¹, Aaron O'Dea^{32,33}, Daniel Pauly³⁴, Péter Szabó^{35,36}, Jimena Torres^{37,38}, John Waldman³⁹, Catherine West⁴⁰, Liqiang Xu⁴¹, Hirokazu Yasuoka⁴², Philine S. E. zu Ermgassen⁴³, Kyle S. Van Houtan^{44,#}

¹Department of History, University of Victoria, Victoria, BC V8W 2Y2, Canada

Full author addresses are given in the Appendix

ABSTRACT: Historical ecology draws on a broad range of information sources and methods to provide insight into ecological and social change, especially over the past ~12 000 yr. While its results are often relevant to conservation and restoration, insights from its diverse disciplines, environments, and geographies have frequently remained siloed or underrepresented, restricting their full potential. Here, scholars and practitioners working in marine, freshwater, and terrestrial environments on 6 continents and various archipelagoes synthesize knowledge from the fields of history, anthropology, paleontology, and ecology with the goal of describing global research priorities for historical ecology to influence conservation. We used a structured decision-making process to identify and address questions in 4 key priority areas: (1) methods and concepts, (2) knowledge co-production and community engagement, (3) policy and management, and (4) climate change impacts. This work highlights the ways that historical ecology has developed and matured in its use of novel information sources, efforts to move beyond extractive research practices and toward knowledge co-production, and application to management challenges including climate change. We demonstrate the ways that this field has brought together researchers across disciplines, connected academics to practitioners, and engaged communities to create and apply knowledge of the past to address the challenges of our shared future.

KEY WORDS: Community engagement · Knowledge co-production · Ecological restoration · Conservation policy · Environmental management · Climate change

1. INTRODUCTION

Historical ecology is a field of research that addresses how human–environment interactions shape ecological change. This rather recent research approach (at least nominally) emphasizes quantifying environmental change and describing the historical

context connecting biophysical and human processes (Szabó & Hédl 2011). Research in historical ecology focuses on the causes and consequences of changes caused by past human actions (Beller et al. 2020), as well as understanding natural variation before and after human intervention (Rick & Lockwood 2013). Although terrestrial ecosystems were initially empha-

*Corresponding author: lorenm@uvic.ca

#These authors contributed equally to this paper

© The authors 2024. Open Access under Creative Commons by Attribution Licence. Use, distribution and reproduction are unrestricted. Authors and original publication must be credited.

sized (Swetnam et al. 1999), marine and freshwater ecosystems later became prominent. Marine historical ecology has pioneered the development of more informed baselines over timescales relevant for management (Pauly 1995) and the documentation of severe impacts to coastal ecosystems due to chronic overexploitation (Jackson et al. 2001).

A central feature of historical ecology is employing a diversity of methods and connecting a range of disciplines—especially history, anthropology, paleontology, and ecology. As a result, historical ecology benefits from source materials that span decades to millennia, and frequently focus on changes from the Holocene to 1950 C.E. (Rick & Lockwood 2013). Historical ecology overlaps in theme with conservation paleobiology, with the latter drawing primarily on fossil and subfossil data and also emphasizing deeper time periods (Rick & Lockwood 2013). Archaeological and paleontological data provide insights from the fossil record, including bones, teeth, shells, and pollen, which can extend from recent to pre-human history (Dillon et al. 2022). Historical ecology also has strong overlap with studies of Indigenous and other local knowledge, and in that context often focuses on the relationship people have built and tended with the environment centered on reciprocal processes (Armstrong et al. 2017). Such local knowledge includes oral histories and community science data streams, which provide observations of ecological change and traditional practices for managing resources (Sullivan et al. 2009, Kittinger et al. 2013, Reif et al. 2021). Finally, historical ecology overlaps with environmental history, including a focus on archival sources, which can range from texts written in medieval times to paintings, newspapers, tax records, menus, as well as records from early natural historians and ecologists who documented their collections systematically (McClenachan et al. 2012). Now a highly transdisciplinary field, historical ecology frequently integrates qualitative and quantitative data (Pandolfi et al. 2003, Lotze et al. 2006, 2011a, Cramer et al. 2021, Paulson et al. 2021) and includes academic researchers and conservation practitioners.

Applications from historical ecology are relevant to conservation and restoration at the species, ecosystem, and landscape scales. Knowledge of multidecadal, centennial, or millennial processes can significantly modify conservation policies based on sub-decadal observations (Lyman 2012, McKechnie et al. 2014, Wolverton et al. 2016). Benchmarks and reference points, such as species recovery goals and other ecological restoration targets that are informed by

longer-term records, can help to avoid the shifting baselines syndrome (Pauly 1995), or the gradual erosion of knowledge about the potential for ecosystem productivity or resilience. Knowledge of past faunal and floral abundances and ecosystem resilience has been used to understand nature-based solutions and develop well-informed conservation plans (Jackson et al. 2001, Balée 2006, Al-Abdulrazzak & Pauly 2017, Nicholson et al. 2024). As a result, historical ecology provides critical long-term context for the current global environmental challenges (Rost 2018, Lotze et al. 2022) and forms a strategic response to initiatives like the United Nations Decade on Ecosystem Restoration (Fischer et al. 2021).

Since early landmark analyses over 25 yr ago (Pauly 1995, Swetnam et al. 1999, Jackson et al. 2001), historical ecology has become a well-recognized field of research that integrates knowledge across disciplines, environments, and geographies. Over this time, multiple initiatives have endeavored to apply historical ecology to conservation, which requires both cross-disciplinary work and engagement with non-academic stakeholders. An increasing focus on diversity, equity, and inclusion, representation, and decolonization has also encouraged researchers to evaluate past methods and interpretations more critically. For historical ecology, this has meant reconciling extractive research practices with the need for co-production of knowledge and grappling with diverse views on historical change and management derived from Indigenous and other local knowledge-holders.

Given all of these developments, there is a strategic need to bring diverse perspectives together to evaluate the state of historical ecology, establish future priorities, and build effective conservation strategies in light of climate change and other anthropogenic perturbations. In this study, we review historical ecology as it applies to conservation, targeting both those already working in historical ecology, but also people that may not be familiar with the goals and aims of historical ecology. We gathered leading and emerging world experts to identify and define the top priorities that will help shape future progress in the field and build on previous syntheses by focusing specifically on conservation (e.g. Balée 2006, Szabó & Hédl 2011, Rick & Lockwood 2013, Armstrong et al. 2017, Beller et al. 2020, Crumley 2021). What have we learned? What persistent problems remain? Where are emerging opportunities? As our objective is to integrate a range of voices, we prioritized expert representation from diverse demographics, quantitative and qualitative perspectives, and alternative forms of knowledge and worldviews.

2. METHODS

We followed the form and process of a series of conservation-themed reviews focused on megafauna conservation published in this journal over the past decade (Lewison et al. 2012, Rees et al. 2016, Jorgensen et al. 2022). Specifically, a steering committee (K. S. Van Houtan, L. McClenachan, T. Rick, R. H. Thurstan, and A. Trant) established the scope of the study and created an initial list of international experts from terrestrial and marine ecology, history, archaeology, anthropology, and paleoecology. We then invited coauthors via email to join the project, using chain referral or snowballing (Biernacki & Waldorf 1981, Van Houtan & Kittinger 2014) to identify potential additional contributors, with a particular focus on increasing diverse participation and global representation.

Next, we employed a structured Delphi process (Linstone & Turoff 1975) to prioritize and select content topics for the review. For the latter, each coauthor submitted up to 4 questions they considered essential to applying the discipline of historical ecology to conservation. We grouped 93 such submissions into 4 category themes. The steering committee excluded questions that were too narrow in scope or not directly related to historical ecology and merged the remaining related submissions into 26 questions. Coauthor voting ranked and focused this list, with questions and category themes further merged by the steering committee where topics overlapped. Coauthors self-selected questions they might answer, with the steering committee then assigning 2 co-authors to each question. Contributors subsequently jointly drafted a response that aimed to summarize the scholarly advances to date, articulate remaining unresolved issues, and identify potential frontiers for future study. All responses were then made available to the full co-author list for comment and review.

To document a baseline of scholarly activity, we queried 3 academic search databases for historical ecology publications over time. To accommodate for differences in database indexing methodology, open-access, search algorithms, disciplinary focus, and literary sources (Martín-Martín et al. 2018), we queried 3 independent sources: Google Scholar (scholar.google.com/), Scopus (www.scopus.com/), and CORE (core.ac.uk/). From 1970 through 2022, we tabulated the number of annual publications containing either of the exact phrases 'historical ecology' and 'historical ecological,' separated by the Boolean operator 'OR'. We report the raw values from each search engine and perform summary statistics on their

ensemble average. Due to the broadly defined, interdisciplinary nature of historical ecology, this search is a broad tool, is inherently imperfect, and does not capture all publications that may be considered to fall within the field.

As observational and cultural norms are framed and legitimized by historical contexts and social settings (Taylor 1992), throughout the process of this project we prioritized the representation of historically underrepresented communities — diverse voices and demographics, qualitative perspectives, and Indigenous knowledge. We particularly sought contributions from women, early-career scholars, and researchers from the Global South. To monitor and self-report our progress, we distributed a survey (available in the Supplement at www.int-res.com/articles/suppl/n054/p285_supp.pdf) to all project coauthors and present the results from all respondents ($n = 37$).

A third-party repository at GitHub (bit.ly/477TePD) provides the data and scripts used in this study. All survey data were anonymized to remove any personally identifying information, and all visualizations were made in the R computing environment, v. 2022.07.2, Build 576 (R Core Team 2023).

3. RESULTS

We defined 18 questions as essential research priorities for the continued application of historical ecology to conservation, which were split into 4 subject groups: (1) methods and concepts, (2) knowledge co-production and community engagement, (3) policy and management, and (4) climate change (Box 1). Question responses aim to summarize the current state of knowledge, emerging opportunities, and recommendations for future progress. Many of the questions that emerged from our analysis focus on issues of colonialism and Indigenous knowledge. These topics are central to current discussions in historical ecology, but we recognize that not all areas of the world followed the same trajectories as regions with significant settler colonial histories. However, colonialism and associated extractive enterprises permeate all regions of the globe and as such, these issues are broadly relevant to the field of historical ecology.

In our review of historical ecology research, the Google Scholar, Scopus, and CORE search platforms identified an average of 17 374 publications containing 'historical ecology' OR 'historical ecological' from 1970 to 2022. From 1970 to 1989, there was comparatively minor activity, averaging 25 or fewer publi-

i. Methods and Concepts

- Q1. What do qualitative and quantitative approaches contribute to understanding long-term change?
 Q2. How do historical data help understand long-term ecological change?
 Q3. How can diverse data types that span a range of time scales be meaningfully integrated?
 Q4. How can historical ecology leverage quantitative methods, big data, and machine learning?
 Q5. How does bias affect the distribution of knowledge in historical ecology?

ii. Knowledge Co-production and Community Engagement

- Q6. What are best practices for knowledge co-production and community engagement for historical ecology?
 Q7. How can natural history museums expand the relevance of historical ecology?
 Q8. How can historical ecology encourage community participation in cultural heritage preservation?
 Q9. What are equitable approaches to knowledge co-production with Indigenous communities?

iii. Policy and Management

- Q10. In what specific policy and decision-making contexts is historical ecology most relevant?
 Q11. How do we select baseline dates for use in conservation?
 Q12. How can conservation baselines consider complex linkages between people and nature?
 Q13. How did past Indigenous management result in outcomes providing insight today?
 Q14. What unique challenges and opportunities exist for historical ecology in data-poor or colonized countries?

iv. Climate Change

- Q15. What sources are available to document past climate drivers and impacts?
 Q16. How can knowledge of ecological and societal responses to past warming inform conservation and adaptation in the face of future climate change?
 Q17. How can management reference historical climate conditions, given that there may be multiple baselines?
 Q18. How may historical knowledge be relevant for future ecosystem states that have no precedent?

Box 1. Global research priorities for historical ecology to inform conservation. The final list of 18 questions are grouped into 4 categories that we address in this review

cations annually (Fig. 1). The period 1990–2016 shows a rapid annual rate of increase ($r = 0.114$), followed by flat production ($r = 0.001$) thereafter. Though beyond the scope of the present study, the reasons for the recent production plateau may reflect the diversification of terminology into more specialized topics and phrases (e.g. Dillon et al. 2022). The 2017–2022 trend does not match the increasing research production seen in the environmental sciences (Davison et al. 2021) over a similar period, and is unlikely related to the decline in research production resulting from the COVID-19 pandemic (Riccaboni & Verginer 2022), whose impact began in late 2020. Further bibliometric or scientometric research in this area may produce additional important insights.

Our author group represents 6 continents, 12 primary languages, diverse career stages, and institutional sectors including academic, non-profit, government, and museum (Fig. 2). Despite this broad representation, many of this study's contributors self-identify as

mid-career academic ecologists working in the USA or Canada, who are white and primarily speak English. While contributions across disciplines, geographic regions, ethnicities, genders and languages is more diverse, significant progress and opportunity remains to achieve greater representation, inclusion, and equity.

3.1. Methods and concepts

- Q1. What do qualitative and quantitative approaches contribute to understanding long-term change?

Both quantitative and qualitative methods are essential to understanding long-term ecological change. Qualitative approaches can be used to describe human knowledge, perceptions, and decisions, while quantitative data can identify ecological patterns and processes (Clavero 2016). For example, qualitative ethno-




Fig. 1. Annual changes in historical ecology research publications from 1970 to 2022. A query for the whole phrases 'historical ecology' OR 'historical ecological' from the Google Scholar, Scopus, and CORE databases returned an ensemble average of 17 374 publications during the 52 yr record. These sources indicate (a) slow growth in scholarly production from 1970 to 1989, (b) a sharp rise from 1990 to 2016 (annual increase 11.4%), followed by (c) relatively flat production from 2017 to 2022 (0.1%). The most recent pattern (c) does not appear to reflect either the broad trends in research production or impacts arising from the COVID-19 pandemic

graphic and quantitative archaeological data on turtle consumption in Polynesia reveal distinct cultural patterns that contributed to different rates of decline over millennia (Allen 2007). Likewise, combining qualitative archival sources and quantitative DNA analysis have documented ancient introductions of non-native species, revealing long-term human influence on biodiversity (Forcina et al. 2015, Clavero et al. 2016).

Working across disciplines can produce new methods that bring analytical weight and narrative power to our understanding of long-term ecosystem change. In the Florida Keys (USA), for example, historical newspapers and photographs were used to quantify an order of magnitude decline in the size of recreational fish caught over 50 yr (McClenachan 2009a,b), while a range of archival sources were synthesized to describe the cultural and political forces motivating this overfishing (Alagona et al. 2023). In Baja California, Mexico, intergenerational knowledge of fishers, archival records, and archaeological data were incorporated to show long-term population trends from ~12 000 yr ago to the 1960s for green turtles *Chelonia mydas*, demonstrating sustainability until they became fungible or commercial goods (Early-Capistrán et al. 2018, 2020).

While not unique to historical ecology, the integration of qualitative and quantitative information is particularly relevant because there is often an expectation of quantification within ecological analyses of change, and in many instances, qualitative data are the only available information over long time frames. In such cases, qualitative observations (such as categorical or ordinal data) can be converted to semi-quantitative metrics including rankings of ecosystem state based on qualitative rules (Pandolfi et al. 2003) and binary measures such as presence–absence or dominance versus non-dominance of taxa (Cramer et al. 2021). Such methods have facilitated global analyses of ecosystem change over millennia and across ecosystems (Pandolfi et al. 2003, Lotze et al. 2006). Similarly, the use of archival data in ecological models reveals patterns and consequences of population extinctions (McClenachan & Cooper 2008), past food web dynamics and trophic structures (Lotze et al. 2011a), and historical habitat distributions that challenged established ideas on the distribution of tree species (Szabó et al. 2017).

In this effort, researchers must understand and employ best practices from specific techniques, fields, and subdisciplines. For example, a critical examination (and sometimes reformulation or transformation) of historical sources to be used under ecological analytical frameworks is necessary, and researchers endeavoring to use local ecological knowledge should be aware of ethical and cultural considerations inherent in these approaches (Szabó & Hédl 2011, Pooley 2013, McClenachan et al. 2015).

Q2. How do historical data help understand long-term ecological change?

Historical ecology has commonly focused on local changes in individual taxa or communities over past decades, centuries, and millennia driven by human activities (e.g. exploitation, habitat alteration, pollution, invasive species) and sometimes natural environmental fluctuations (Lotze & McClenachan 2013, Rick & Lockwood 2013). From the scale of species to landscapes, historical ecology is important for documenting past occurrences, abundance, distributions, demographics, habitat usage, and species interactions. For example, surveillance satellite data (Munteanu et al. 2020, Rizayeva et al. 2023), historical maps (Fuchs et al. 2015, Munteanu et al. 2015), and aerial photography (Lydersen & Collins 2018) can be used to reconstruct ecosystem structure and species habitat use at regional to global scales extending cen-




Fig. 2. Self-reported perspectives from the contributing authors of this study. Authors' responses to survey questions (questions available in the Supplement at www.int-res.com/articles/suppl/n054p285_supp.pdf) on (a–c) professional affiliation and approach, (d,e) geographic domain, (f–h) demographics, and (i,j) language use. Despite significant representation beyond these categories, most authors of this study are white, English-speaking, mid-career quantitative marine ecologists who conduct research at universities in the USA or Canada

turies in the past. Natural archives (Wolfe et al. 2013, Xu et al. 2016) can be used to quantify ecosystem productivity, community structure, and population sizes at millennial time scales via a large number of organic and inorganic proxies, such as pollen, bio-elements,

stable isotopes, and ancient DNA (Emslie 2021, Palli et al. 2023).

When using historical information to understand ecological change, several challenges persist. Ecosystem variation over decadal to millennial scales is high

(Duda et al. 2023), and available historical sources may miss the full range of variation in ecosystems (Soga & Gaston 2018). Discriminating the effects of biophysical variability and anthropogenic activity remains difficult even if the range of variation is known (Swanson et al. 2009, Armstrong et al. 2017). Finally, when used in isolation, individual historical data sources may be difficult to validate, leading to interpretive errors (Stahl 2018). Several solutions exist. Integrating historical species records, modern observations, and modeling approaches, for example, allows for long-term range and population dynamic reconstructions. Developing more refined tools for integrating these different data sets should be prioritized. For example, specific biomarkers may lead to unambiguous conclusions on human- versus environment-driven changes, and more precise age determinations would shed light on the past at finer timescales, avoiding dating-induced biases (Bluhm & Surovell 2019, Zimmerman & Wahl 2020). Importantly, data on various scales and from different disciplines can help to validate results: population dynamics or vegetation coverage from pollen records can be spatialized using historical maps and remote sensing, and ecosystem dynamics from maps or aerial photographs can be validated using similarly timed literature entries, species records, or biomarkers.

Q3. How can diverse data types that span a range of time scales be meaningfully integrated?

In integrating diverse data, historical ecology researchers confront issues regarding the scale, resolution, and format of data. These can be addressed by standardizing metrics of interest (Lotze et al. 2011b). Oral history, archival, and fossil/subfossil data have been integrated to reconstruct timelines of ecological change, relative abundance of key taxa, and fisheries catch over thousands of years (Lotze et al. 2006, Kittinger et al. 2015, Early-Capistrán et al. 2018). In doing so, calibration across diverse data sets has been achieved by assessing the correlation between metrics, such as the accumulation of shark dermal denticles preserved in marine sediments and underwater surveys of modern shark abundance (Dillon et al. 2020), hunting rates for dugongs from archaeological middens and modern conservation assessments (McNiven & Bedingfield 2008), and information from oral history and commercial fisheries records (Sáenz-Arroyo & Revollo-Fernández 2016).

Several technological and methodological advances are accelerating the integration of data types across

timescales. Increases in instrumentation and artificial intelligence capabilities are increasing the capacity to analyze ancient and environmental DNA, and isotopic, elemental, and histologic data that can be collected in the same manner across varying sources of data (paleoecological, archaeological, collections-based, and modern) (Kidwell 2015). High-precision chronologies from natural archives are available via radiometric dating, providing similar temporal resolution across historical and modern data (Clark et al. 2014). Several approaches would improve the capacity to integrate diverse data to understand long-term change. Using historical data to validate hindcasting based on modern data can help to understand ecological dynamics, such as if past impacts of climate change on species distributions were similar to those observed today (Dietl & Flessa 2011). Improved data integration may also arise from collaborating across disciplines to create standardized methods and deriving conversion factors for ecological metrics (Miller et al. 2019, 2020) by designing studies that allow for statistical comparisons of historical and modern data.

Q4. How can historical ecology leverage quantitative methods, big data, and machine learning?

The large-scale acquisition, storage, and analysis of ecological and related information provides conservation science with unprecedented resources and opportunities (Tuia et al. 2022). Researchers and institutions are responding by embracing new technology and methods in cloud storage and computing, artificial intelligence, automated systems, '-omics,' and community science (NOAA 2020). Similar initiatives are helping to increase the efficiency and productivity of historical ecology research programs, where the extraction and curation of historical data streams are often manual and logistically intensive. For decades, the humanities have focused on digitizing, processing, and publishing relevant historical texts, images, and cultural ephemera (Galla 2009, Van Houtan et al. 2013). Further enhancing accessibility and linking those efforts to the goals of historical ecology, such as the open publication of papers and data sets resulting from The History of Marine Animal Populations Project and Oceans Past Initiative (e.g. Starkey et al. 2012, Polónia 2020), will be fruitful. When standardized and attributed with metadata, open-access clearinghouses of historical data may also further advance the field in the mold of more established (Ratnasingham & Hebert 2007, Beck et al. 2014) and growing

disciplines (Jung et al. 2018, Miller et al. 2022). In addition, techniques to advance automation and robotics in -omics methods (Sinclair et al. 1998, Searcy et al. 2022) will improve the efficiency of data acquisition and analysis across a range of historical ecology topic areas.

With new data come new analytical methods, such as artificial intelligence and machine learning methods. In ecology, these approaches have been used in remote sensing, genetics, climate, and biotelemetry, often by incorporating autonomous sensor networks into research workflows (Tuia et al. 2022). Applying machine learning to qualitative data such as text (e.g. natural language processing and large language models) is rapidly advancing (Shen et al. 2023), with significant potential for conservation (Van Houtan et al. 2020). Such approaches complement traditional historical ecology methods, where expert-trained data sets and advanced algorithms work together to automate learning from historical texts (Davidson et al. 2019) in multiple languages. Beyond removing barriers to historical data, adopting big data

and artificial intelligence in historical ecology may help reduce geographic bias (see Q5 and Thurstan 2022), while increasing access and cultural representation (see Fig. 3, Q8–9). The reduction of such biases also requires addressing existing structural inequalities, for example, access and expertise gaps arising from the concentration of high-powered technological infrastructure in highly developed nations (Obermeyer et al. 2019, Jindal 2022). Assuming such inequalities are addressed, collectively these outcomes will advance a core goal of historical ecology—to increase knowledge for conservation practitioners across contexts and solutions.

Q5. How does bias affect the distribution of knowledge in historical ecology?

The availability of historical ecology data is affected by biases across space, taxonomy, and time. Most fundamentally, historical ecology has, like other Western scientific endeavors, been affected by colonial power




Fig. 3. The Florida Museum of Natural History's Randell Research Center and Calusa Heritage Trail. Located at the Pineland Site Complex on Pine Island, Florida, the museum center is explicitly dedicated to the research, preservation, and public education about the archaeology and historical ecology of the Calusa and Pine Island Sound estuary. (a) Welcome sign clearly naming archaeology, history, and ecology as the cornerstones of the center and trail. (b) Center operations manager and (c,d) museum educators leading Title I School fourth-grade educational programs across preserved Calusa shell mounds, middens, and canals. (e) Classroom display with replica artifacts for public education about estuarine historical ecology and 2000 yr of fishing. Photos provided by Annisa Karim and Charles O'Connor and used with permission

structures, which results in knowledge biases (Fernando 2022). For example, colonial administrations destroyed historical sources, while also producing documents that can help understand the historical ecology of many colonized regions. In using such sources, researchers must assess and account for those biases (McClenachan et al. 2015). Likewise, archaeological sources may contain biases that affect their use, and assessments of the so called 'cultural filter,' or the ways in which objects are biased by human actions such as transport, are needed (Peacock et al. 2012). Additionally, there are socio-economic and geographic biases in analyses of available sources (Trisos et al. 2021, Fernando 2022, Raja et al. 2022). Roughly 90% of historical ecology studies derive from North America and Europe (Beller et al. 2020), for example. Furthermore, biological and cultural objects held in museums or private collections, trapped behind paywalls, or written in languages not spoken by local communities, present significant barriers to equitable access (Meagher 2021).

These cultural biases result in an unequal distribution of records relevant for historical ecology, such as an overrepresentation of studies on environmental change and biodiversity in human population centers across the Global North (Gagné et al. 2020, Hughes et al. 2021), with large data gaps in the biodiversity-rich tropics (Felton et al. 2009, Davison et al. 2021, Spalding et al. 2023). Cultural biases for charismatic and commercially valuable species drive taxonomic biases regardless of time and location (Davison et al. 2021, Daru & Rodriguez 2023), while available data are also region- and taxon-specific and accompanied by taphonomic biases (Barnosky et al. 2017). Temporal biases emerge from technological advancement, accessibility, research interests, and socioeconomic factors changing over time (Stropp et al. 2016), and the tendency to collect historical sources and ecological data based on convenience, such as adjacent to research centers (Davis 2018), perpetuates global biases despite the growth of community science records in recent decades (Sullivan et al. 2009, Hughes et al. 2021, Daru & Rodriguez 2023). Biased ecological data distort our understanding of biodiversity gradients, species distributions, and predictions of their responses to human-driven change (Hortal et al. 2008, Gagné et al. 2020, Daru & Rodriguez 2023). Solutions include developing mechanistic models of ecological distributions (Reygondeau et al. 2020), addressing sources of cultural biases, and amplifying and broadening knowledge systems from developing countries, marginalized communities, and Indigenous and other local knowledge co-production (see Q6, Q8–9).

3.2. Knowledge co-production and community engagement

Q6. What are best practices for knowledge co-production and community engagement for historical ecology?

Community engagement in historical ecology ranges from knowledge dissemination to the public in the context of natural history museums (see Q7) to the co-production of research priorities with Indigenous knowledge holders (see Q9). Given this wide-ranging and often complex social setting, there is no single set of best practices for knowledge co-production and community engagement for historical ecology. Yet, several principles and practices are relevant in many contexts. First, accessibility of information is needed, because despite the transdisciplinary nature of historical ecology, information sharing remains restricted by factors including the disciplinary siloing of knowledge (Carmenta & Vira 2018), competition among researchers, and the privileging of specific types of knowledge and holders (see Q5) (Sidik 2022). Platforms for open-source data sharing (see Q4) and guidelines for ethical best practice (Wilkinson et al. 2016, Crystal-Ornelas et al. 2022) facilitate information sharing to meet global conservation challenges. Likewise, knowledge may be better shared through audio-visual tools such as infographics, videos, and the classroom, and relevant materials should be communicated in locally spoken languages (ISE 2006, Lawson et al. 2021).

While improved communication training and the implementation of wider dissemination strategies can enhance information sharing in historical ecology, they often remain unidirectional as they may host data 'from' researchers 'for' others to use. Instead, more fundamental changes are needed to ensure equitable representation of historically marginalized histories, voices, and knowledge systems. For example, 'community peer review' is a mechanism which supports research and knowledge co-production led by community groups. Community peer review includes steps to ensure that research outputs (papers, data, etc.) are accurate and consistent with community standards for data sharing (Liboiron 2021). This process helps to guard against parachute science or tokenism, where knowledge is extracted and misapplied without consent (de Vos & Schwartz 2022). Appreciating the deep entrenchment of local knowledge (Klenk et al. 2017) and recognition of oral traditions and knowledge systems (Reid et al. 2021) is essential in this process, as is taking steps to commu-

nicate the differences across knowledge forms and examine how such differences may influence their acceptance, application, and interpretation. Moreover, ethical engagement and community self-determination must be the foundation for collaboration (ISE 2006, Mistry & Berardi 2016, Schroeder et al. 2019).

Emerging models of information sharing could also be more broadly incorporated into historical ecology. For example, collaborative research learning networks (RLNs) promote more inclusive models of knowledge co-production and information sharing (Dalton et al. 2020), and are emerging as alternatives to traditional institutional and academic networks. RLNs, such as the current UN Ocean Decade (Singh et al. 2021), promote formal working groups with wide representation that aim to collectively build a comprehensive and accessible knowledge base while fostering collaboration and inclusivity (Gerhardinger et al. 2020). Blending RLNs with enhanced media and information literacy could further facilitate knowledge co-production and sharing, empower individuals and communities, and more effectively navigate complex environmental issues (Gerhardinger et al. 2020).

Historical ecology increasingly recognizes the pivotal role that communities play in the framing and prioritization of research questions, data curation, and conservation efforts (Evans et al. 2008, Greer 2010, Shackeroff et al. 2011). The focus of historical ecology on place and potential for powerful imagery can act as a catalyst to break down barriers to community participation (Astudillo et al. 2023). Additional expertise and training in inclusive community engagement, such as participatory or consensus-based techniques (Wells et al. 2019), may be required. Such relationship building requires significant time and resources from the beginning of research, and extends beyond typical project timelines (Mace 2014, Morgan et al. 2023).

Q7. How can natural history museums expand the relevance of historical ecology?

Although rarely nominally referenced, natural history museums have long featured some of the most prominent tenets of historical ecology in their research and public engagement. These include museum collections, exhibits, research, and educational outreach on climate and ecosystem change, evolution and biodiversity loss, and the dynamics of human–environment relationships. Such themes are often presented across multiple time scales represented by

paleontological, archaeological, and historical natural history collections. Recently, however, the historical and societal relevance of natural history museums as collection repositories and bastions of public education have been critiqued (Dorfman 2017, Reid & Vail 2019). Natural history museums' long-professed interpretive neutrality when exhibiting natural and cultural history is now increasingly recognized as problematic and perhaps a disingenuous representation of the featured ecosystems and people (Evans et al. 2020, Shiraiwa & Zabalueva 2021). Rather, 'museum activism,' has emerged as the way of the future for relevant natural history museum practice and exhibits. This approach centers purposeful community engagement and co-production in research, exhibits, public education, and solutions to pressing societal challenges (Robinson 2017, Drotner et al. 2019, Janes & Sandell 2019, Sutton 2020).

It is within this spirit that natural history museums are well-positioned to expand historical ecology from the status quo, which is more of an underlying academic framework. Beyond increasing access to and diagnostic research on collections (see Q4), this transformation may move to more publicly recognized, community-accessible, applied science focused on understanding how the past can help address Anthropocene challenges (Miller et al. 2019). Doing so may help communicate how linked ecological and cultural histories have together shaped socioecological systems through time (Fig. 3). This work requires unpacking colonial structures through consultation and collaboration with the communities whose histories, legacy environments, and continuing biocultural heritage are represented in collections and exhibits. Engagement with local biological and cultural heritage stakeholders is essential. Natural history museums are often the public face of historical ecology, and bear responsibility for shaping and demonstrating its societal, scientific, and conservation relevance. At the same time, these principles also apply to local education centers and classroom settings, in particular in areas of the world where natural history museums lack access.

Q8. How can historical ecology encourage community participation in cultural heritage preservation?

Like natural history museums (see Q7), the preservation of cultural heritage is not always associated with historical ecology, but it can be a powerful mechanism for engaging communities. Cultural heritage is

both tangible (e.g. cultural landscapes) and intangible (e.g. traditions) and intertwined with people's interactions with local environments (Silverman & Ruggles 2009, Hassan 2014). Historical ecology research can help identify cultural heritage and its significance to communities in time (Crumley 2021, Astudillo et al. 2023). Leveraging community knowledge via historical ecology research applying community-based approaches can empower communities to protect cultural heritage by engaging with drivers of heritage decline (Piñar & Sterflinger 2021, Curran & Zimmermann 2022), reconnecting with and redefining their heritage (Astudillo et al. 2023), reframing dominant discourses (Hill et al. 2011, Koslov 2016), and generating information for consensus-driven policy and practice (Hill et al. 2011, Armstrong et al. 2017). In aiding understanding of past environmental changes and their drivers, historical ecology can illuminate how heritage practices interacted with and influenced local environments. For example, the rediscovery of lost 'dreg' songs from the south coast of the Firth of Forth, which used a specific song pattern to maximize the efficiency of the oyster dredges in this particular environment, and which went extinct with the fishery, became the focus of a community effort to reintroduce these songs and subsequently illuminated the loss of the native oyster in the Forth (Bishop et al. 2013). Such understanding helps communities to reframe their priorities and build future visions that facilitate community adaptation and transformation in the face of environmental change and heritage loss (DeSilvey & Harrison 2020, Hassan 2021).

Q9. What are equitable approaches to knowledge co-production with Indigenous communities?

Indigenous and other local knowledge is valuable for generating a more holistic understanding of ecosystems and the human–environment relationship through time, which is central to historical ecology (Spalding et al. 2023). Combining Indigenous and scientific knowledge as complementary approaches (McKeon 2012, Hall et al. 2015, Marshall et al. 2015, Abu et al. 2020) is referred to as '2-eyed seeing' (Bartlett et al. 2012); this methodology values and amplifies the intellectual tool box that Indigenous peoples carry, and draws from daily activities and cross-generational factors when engaging in socio-cultural and ecosystem-based activities (see also 'braiding knowledge'; Atalay 2020). Central to this methodology is ensuring that the research benefits all

(Bartlett et al. 2012), and does not exploit Indigenous people.

A persistent challenge facing conservation practitioners is how to work with Indigenous and other local knowledge respectfully and ethically throughout the processes of conservation. Two related models for partnership with Indigenous and other local communities include equitable exchange (Harris et al. 2021) and 'boundary spanners' (Hatch et al. 2023). Equitable exchange focuses on 'currencies' within different communities, and proposes a framework for defining these currencies prior to formal partnerships with the goal of equal and self-defined benefit. The success of many mainstream-community partnerships is frequently dependent on a boundary spanner—a person or small team that is fluent in both community protocol and mainstream science. Unlike the concept of 'informant' in anthropological research contexts, the boundary spanner is often a member of both communities, and helps to ensure that equitable exchange occurs. Future progress may also be found in the Indigenous 'land back' and 'water back' movements (Fisk et al. 2021, Gray 2022, Leonard et al. 2023). These efforts are rooted in Indigenous peoples' sovereignty over their traditional territories, and focus on the use of local ecological knowledge to support Indigenous conservation methods (Simpson 2017).

Equitable approaches to knowledge co-production also integrate ethical data management, and acknowledge Indigenous authority over intellectual property rights. Methodologies that ensure data sovereignty and protect intellectual property rights are essential to knowledge sharing and co-production relationships. Strategies within this realm include community-controlled information databases such as Mukurtu (<https://mukurtu.org>), protocols for Indigenous intellectual property use, and memoranda of agreements (Buck & Hamilton 2011, Malsale et al. 2018, Carroll et al. 2020).

3.3. Policy and management

Q10. In what specific policy and decision-making contexts is historical ecology most relevant?

Historical ecology is relevant for policy and decision-making involving the management of species, populations, and ecosystems. Historical research has helped resolve the biogeography of priority species (Clavero et al. 2016), justify source populations for species reintroductions (Gessner et al. 2011), and

challenge modern ideas of 'natural' ecosystem states (Jackson 1997, Szabó et al. 2017). One such case involves the 'American' Atlantic sturgeon *Acipenser oxyrinchus* and European sturgeon *A. sturio*. These species diverged at least 15 million years ago (Ludwig et al. 2002), and it was thought that Atlantic sturgeon did not exist in European waters. However, historical research documented that humans extirpated Atlantic sturgeon from European waters in the 1600s (Desse-Berset 2011), which supported enhancing Baltic sturgeon populations with the more abundant Atlantic sturgeon instead of the Critically Endangered European sturgeon (Gessner et al. 2011).

Historical ecology can help communities converge on shared goals for restoration (see Q8). In South Australia, for example, a multidisciplinary network of scientists, practitioners, and managers was successful in reaching out to a diversity of stakeholders to communicate the past (and potential future) social, ecological, and economic significance of oyster (*Ostrea angasi*) reefs. This helped to build political support for the establishment of a large-scale restoration initiative and a nationwide 'reef-building' agenda (McAfee et al. 2022). Pooled data from explorer histories, fisheries catch, archaeological records, and sediment cores contributed to a Critically Endangered listing of Australian oyster reefs on the IUCN Red List of Ecosystems (Gillies et al. 2020). Here, historical ecology motivated and legitimized government intervention, including customizing restoration for the current environment and paired social benefits (McAfee et al. 2021).

Finally, historical approaches are necessary to historicize and critique ideas like management itself, as Western scientific management exists within a specific set of cultural frameworks and biases. For example, in southwestern Ethiopia, fire plays a crucial role in the conservation of African savannas and as a tool for nomadic pastoralists to create pastures. The establishment of a national park in 1980 led to a fire ban for local communities, particularly the Mursi people (Turton 2011), as high-frequency fires were thought to cause bush encroachment and to reduce grass availability for wild ungulates. However, a combination of paleoecological indicators (Gil-Romera et al. 2010), Mursi oral histories (Gil-Romera et al. 2011, Turton 2011), and current-day plant ecology revealed that bush encroachment was related to fire suppression policies (Gil-Romera et al. 2010). This example demonstrates that by adding more voices to the field of historical ecology, it will become more complex, more meaningful, and allow more people to join a respectful community of practice.

Q11. How do we select baseline dates for use in conservation?

Baselines are ubiquitously, though often implicitly, used in conservation endeavors (Alagona et al. 2012). Such reference points from the past simultaneously provide targets to guide management actions and thresholds for assessing recovery progress or status (Grace et al. 2021). Poorly-chosen baselines—whether due to unacknowledged assumptions, cultural biases, or shifting baselines—can have negative consequences, such as the premature removal of protection (e.g. Yellowstone wolves; Sturges 2019) and misleading assessments of population trends (e.g. furbearer hunts, Collins et al. 2020; also see Q13, Q17).

Why are baselines selected, and by whom? The IUCN Green Status of Species (IUCN Species Survival Commission 2021), a recovery assessment, identifies species-relevant baselines between 1500 and 1950, recognizing that human impacts occurred outside of that period, but asserting that earlier baselines are too removed from the present (Stephenson et al. 2019). Policy baselines are often set later, within the past 50 yr (Burns et al. 2023). Choice of baselines is also vulnerable to cultural biases that render them value-laden. The initial objective of US National Parks, for example, was to maintain or re-create ecosystems 'in the condition that prevailed when the area was first visited by the white man' (Leopold et al. 1963, p. 3). Recognizing the biases inherent in this objective, it has since been updated to the more open-ended directive 'to preserve ecological integrity and cultural and historical authenticity.'

Another consideration is whether reference periods are selected *a priori* or emerge from a synthesis of historical data. *A priori* baselines include inflection points in humanity's relationship with the environment and each other, as occurred during the climatic shifts and megafaunal extinctions of the Late Pleistocene/Holocene transition (~11 000 yr ago and earlier), the onset of European colonization and the widespread dislocation of Indigenous communities (ca. 1500), in the Industrial Revolution (beginning ca. 1750), and the Great Acceleration (ca. 1950) (Ellis et al. 2016). However, not all species were affected equally by these shifts in human activity, so synthesis of species-specific historical data may be necessary to choose a meaningful baseline. It is also important to remember that not all conservation goals are backward-looking. In some cases, historical baselines are put aside or used as a menu of options when helping species, communities, and ecosystems adapt to a changing

world (Barnosky et al. 2017, Coleman et al. 2020, Mychajliw et al. 2022).

Q12. How can conservation baselines consider complex linkages between people and nature?

One important recent shift in historical ecology is from a focus on informing conservation baselines with information before human impact (i.e. so-called 'pristine' conditions), to one that emphasizes the role of human societies in shaping both lasting ecological patterns and socially valuable outcomes. Historical ecologists have described human agency in landscape formation, even in seemingly pristine forests, where intermediate disturbances by people sometimes enhanced the biodiversity in landscapes (Denevan 1992, Balée 2006, 2013, Balée & Erickson 2006, Yasuoka 2013, Clement et al. 2015, Levis et al. 2017, Avtzis et al. 2018). People developed complex socio-ecological arrangements all over the world, many of which have been disrupted over past centuries by growth-centered capitalist (Rose 2004) and modernizing forces — including state-sponsored conservation, which is often based on the human–nature dichotomy (See also Q10) (West et al. 2006, Fletcher et al. 2021). Many protected areas overlap lands where Indigenous peoples and local communities historically lived (Garnett et al. 2018). Therefore, using 'pristine environments' as benchmarks for conservation baselines, while untenable as it is often embedded within colonial discourses (Early-Capistrán et al. 2018), causes normative, scientific, and practical problems (Hilding-Rydevik et al. 2017). For example, historical analyses are challenging views on management of keystone species, such as sea otters in British Columbia and Alaska, as evidence emerges that Indigenous communities managed otter populations in order to enhance shellfish productivity (Salomon et al. 2015). Coupled with justice-informed research, this work demonstrates that the recovery goals established for otters may benefit non-Indigenous stakeholders and harm historically disadvantaged Indigenous fishers (Levine et al. 2017). When discussing ecosystem restoration, historical ecology is increasingly demonstrating the need to acknowledge the ecological roles of people in ecosystem function (Bliege Bird & Nimmo 2018). Conservation can in fact become more effective by reinforcing the role, capacity, and rights of Indigenous peoples and local communities (Dawson et al. 2021). Such shifts in perspective can be controversial, as it challenges established knowledge priorities within conservation biology.

Baselines are socially constructed and thus influenced by political agendas, economic realities, preconceived ideas, and socio-cultural understandings of human impacts on nature (Hilding-Rydevik et al. 2017). Therefore, to share more appropriate baselines and collaborate with local peoples in a dynamic world, knowing the past, as well as ongoing, human–landscape interactions, even if imperfect, will be an indispensable guide (Alagona et al. 2012). Future work in historical ecology that combines scientific and Indigenous knowledge to focus on the complex material and cultural linkages between people and nature will help scholars, decision-makers, and community members articulate and attain support for ambitious conservation goals that seek to repair and conserve nature while respecting local knowledge and traditions and promoting environmental justice.

Q13. How did past Indigenous management result in outcomes providing insight today?

For millennia, Indigenous peoples throughout the world have directly and indirectly shaped ecosystem dynamics, resulting in a continuum of outcomes from enhancement to degradation. Habitat modification by Indigenous peoples includes structural changes, such as building stone walls in the intertidal, as well as changes in ecological processes like fire and propagating and hunting species (Boivin et al. 2016, Ellis et al. 2021, Hoffman et al. 2021). Bringing together historical ecology and Indigenous knowledge offers insight into the scale and scope of past management and stewardship activities. Importantly, these insights can empower Indigenous communities, guide place-based restoration, and help identify conservation priorities.

For example, in forest islands of the Llanos de Moxos, in the Bolivian Amazon, shellfish gatherers visiting seasonally flooded grasslands for more than 10000 years have built-up middens on higher ground, which transformed into islands supporting trees and other dense vegetation (Lombardo et al. 2013). These constructed soils provided nutrient inputs to support intensive traditional agricultural production starting around 5000 yr ago to present day (Lombardo et al. 2020). In the Western Desert of Australia, the Indigenous practice of deploying hunting fires buffers against larger-scale, and increasingly climate-driven, wildfire and creates more diverse landscape mosaics (Bliege Bird et al. 2008, 2016), increasing food and shelter for common wallaroo *Osphranter robustus* (Coddington et al. 2014) and monitor lizards (*Varanus*

spp.) (Bliege Bird et al. 2013), which are critically important food for Indigenous peoples (Bliege Bird & Bird 2021). On North America's Pacific Coast, the construction of clam gardens by coastal Indigenous peoples increased the available niche space for clams and secured a reliable food source for people (Smith et al. 2019, Toniello et al. 2019, Lepofsky et al. 2021). Clam gardens have had persistent indirect effects on soft sediment marine communities (Cox et al. 2019) and increased nearby forest productivity around village sites where clam shells and other materials were deposited after being consumed (Trant et al. 2016). Clam garden restoration is occurring today in many Indigenous territories (Wickham et al. 2022). While most commonly studied in settler colonial regions, similar lessons emerge from research in other parts of the world. For example, coppice management in pre-modern Europe maintained higher plant diversity in lowland forests than that found in unmanaged closed forests (Hédli et al. 2010).

Solutions gleaned from Indigenous stewardship and historical ecology include strategies that increase productivity and crop yields through the addition of soil amendments, enhance habitat conditions for culturally important foods through use of fire, and set conservation and restoration targets (Allen 2007, Evans et al. 2008, Lotze et al. 2011a, Colonese et al. 2023). While all of these practices are critical for Indigenous food security and food sovereignty, they may also shed light on contemporary conservation and restoration strategies more broadly.

Q14. What unique challenges and opportunities exist for historical ecology in data-poor or colonized countries?

Knowledge on the dynamics of populations and ecosystems on centennial and millennial scales has recently increased in low- and middle-income countries in the Global South. The unique challenges that these countries face in engaging with historical ecology research are that most were subjected to European colonial expansion, which led to the abrupt imposition of new elites and their languages and the silencing of the original surviving inhabitants (Palmare et al. 2007). As a result, much of the knowledge of colonized societies, including those with written records (e.g. the Maya, whose books were burned; Christenson 2007) was lost. Thus, while the time horizon that historical ecologists choose to work on tends to depend on the ecosystem they study and on the study's goals and outcomes, those working in coun-

tries that were colonized will be faced with information gaps related to the damage of the linguistic and cultural rupture that colonialism caused. Additional challenges include the maintenance of archives and colonial science which results in knowledge being controlled by outsiders.

Certain countries in the Global South face challenges such as political instability and limited funds, coupled with priority areas for fund allocation that may perceive disciplines related to historical ecology as less strategically aligned with economic development. For example, Brazil, despite being a biodiversity hotspot, has experienced substantial cuts in research and disciplines within the social and humanities realms. This issue is not exclusive to Global South countries, but increasing populism and political polarization have a more pronounced impact on low-income and less literate nations than countries in the Global North (Moutinho 2022). These political challenges underscore the importance of engaging with local communities, private sectors, institutions such as museums (see Q7), and the general public to enhance the visibility of historical ecology, and promote its application and training at both undergraduate and postgraduate levels.

Historical ecology also presents unique opportunities in data-poor contexts, particularly because ecological knowledge in most post-colonial countries is limited by the short duration of continuous instrumental records (e.g. atmospheric conditions) and observation records (e.g. fisheries statistics). In this context, historical, local knowledge, archaeological, and paleo-environmental sources are vital, as they can provide quantitative evidence of ecological status for pre-instrumental periods (see Q4). In addition, such sources can help estimate ecosystem conditions before crucial historical turning points triggered by European contact, such as the establishment of a market economy (e.g. Fossile et al. 2023) or recent policy incentives promoting industrial exploitations (Herbst et al. 2023). The information gained can provide strong arguments for the protection and conservation of ecosystems.

3.4. Climate change

Q15. What sources are available to document past climate drivers and impacts?

Linking historical changes to underlying biophysical drivers at the millennial and basin scale has long been of interest (Swetnam et al. 1999), but progress has been limited, partly because of the mismatch of

temporal and spatial scales and the resolution of available data (Lotze et al. 2011b, 2022, Harnik et al. 2012, Finnegan et al. 2015, 2024, Beller et al. 2020). With increasing coverage of historical, archaeological, and paleoecological data plus advances in modeling for hindcasting and mapping, integrating various data sources can help discern between drivers of change, and inform future projections (Lotze et al. 2019, Sandweiss et al. 2020). For example, in the Northwest Atlantic, future climate change may shift ecosystems towards conditions last seen >5000 yr ago, with warmer waters, red tides, and other algal blooms being much more prevalent than in recent centuries (Lotze et al. 2022). Similarly, synthesizing global paleoecological records during the past 21000 yr illustrates that future warming will cause major changes to terrestrial ecosystem composition and biodiversity (Nolan et al. 2018).

To better resolve long-term (centuries to millennia or more) changes over larger spatial scales, additional historical records from terrestrial and aquatic inshore and offshore sources are available (e.g. marine sediments, mollusk shells, corals, animal bones and teeth, and phytoliths; also see Q2, Q4) (Sherwood et al. 2011, Sibert et al. 2017, Nolan et al. 2018, Lotze et al. 2022). Comparing patterns to local or regional ecological changes, often documented in historical and archaeological sources, will help address foundational questions about environmental drivers of change. For instance, integration of different environmental proxies (pollen from lake sediments, dendrochronology, historical maps, etc.) document the drivers of changing vegetation composition in the Mediterranean over the past millennium, with implications for understanding future vegetation change in an uncertain climate (Palli et al. 2023).

Q16. How can knowledge of ecological and societal responses to past warming inform conservation and adaptation in the face of future climate change?

The structure and composition of ecological communities are, to a large extent, driven by climate at millennial and global scales (Jackson & Erwin 2006, Zhang et al. 2007, Deutsch et al. 2015, Reddin et al. 2022). Historical environmental data are central to resolving models that seek to provide robust predictions of ecological responses to future climate conditions (Crowley & Berner 2001, Lotze et al. 2019, 2022). On land, records of past environmental and ecological dynamics reveal the thermal fragility of high mountain flora that are likely to be driven

extinct along with their endemic fauna as temperatures rise (Flantua et al. 2019). In contrast, lowland tropical forests may be able to tolerate rising temperatures, but only if rainfall persists (Slot & Winter 2018), demonstrating the importance of modeling precipitation patterns as well as temperature to inform conservation. With enough spatial resolution, prior records are well suited to help resolve several key questions facing conservation, including the long-term threat of extinction debt (Jackson & Sax 2010) and the ability (or lack thereof) of ecological communities to persist under changing climates (Staples et al. 2022, Wang et al. 2023).

Due to its inherent extended chronology, historical ecology is uniquely situated to evaluate past societal responses and adaptations to major weather events and climate variability. A more accurate depiction of peoples' past responses to climate change cuts through assumptions and biases of peoples' knowledge, capabilities, success, and failures when mitigating socio-ecological risks from changing weather patterns, sea level, and climate variability (Schaan 2016). For example, people have been shown to exploit the expansion of fire-prone landscapes under changing climates (Hoffman et al. 2021). Mid-Holocene terracing and terraforming of the intertidal across the Pacific Northwest was a highly localized but broadly applied practice for dealing with sea level changes while increasing food production (Lepofsky et al. 2021). Pastoralism has been highly variable and adaptive over space and time, despite (incorrect) assumptions about its heterogeneous practice and regional impacts (e.g. overgrazing) (Boles et al. 2019). Local Indigenous knowledge is particularly valuable in understanding how communities have adapted livelihoods and cultures in response to climate change (Reyes-García 2024). Understanding the ways in which societies adapted to changing climates throughout history provides insights into societal responses to modern warming and can help to inform more equitable resilience planning (Crumley 1994, Degroot et al. 2021, Reeder-Myers et al. 2022).

Q17. How can management reference historical climate conditions, given that there may be multiple baselines?

As world temperatures continue to rise and affect ecosystems, multiple baselines may be beneficial and practical. Historical reference conditions, at least

regionally, can be selected from records of past analogues of current climatic trends, such as the Last Interglacial (~110000 yr ago), the mid-Holocene Climatic Optimum (~8000–4500 yr ago) and the late Holocene Medieval Climate Anomaly (MCA, ~1300–650 yr ago) (Jerardino 2012, Rick et al. 2020). Quantitative data derived from these records can be used to characterize ecosystem functions at that time and socio-environmental responses to climate change, providing insights into current warming trends (Meltzer 1999, Rick et al. 2020). Paired with recent written records of species and habitat occurrences and their use (e.g. fisheries records), these data can establish targets for ecological conservation (Braje et al. 2016). However, few studies have used data on specific past environments for projecting into the future (e.g. Rivera-Collazo & Perdikaris 2023), and while lists of taxonomic abundances are described for past environments, it is not always straightforward how they translate into baselines (see Q11). Therefore, close collaborations of historical ecologists and conservationists are vital for linking these disparate data sets.

Q18. How may historical knowledge be relevant for future ecosystem states that have no precedent?

Historical knowledge has significant value for guiding ecosystem restoration, resource management, and conservation under conditions that are unprecedented or novel in the recent past (Higgs et al. 2014). Historical knowledge in many forms (Higgs et al. 2014) can illustrate past variability to explain ecological legacies, contextualize unprecedented change, and provide a set of possible expectations for and responses to unpredictable climatic and ecological conditions (Grayson 2005, Millar et al. 2007, Crumley 2021).

For example, catastrophic marine heat waves threaten many fisheries, and North Pacific fisheries managers have called for the integration of a historical perspective into management strategies (Barbeaux et al. 2018, 2020). Historical data suggest there have been significant climate-driven periods of change in the Pacific herring *Clupea pallasii*, Pacific cod *Gadus macrocephalus*, and salmon (*Oncorhynchus* spp.) fisheries over millennia (West et al. 2022). Both managers and archaeologists (Barbeaux et al. 2020, West et al. 2022) have drawn on these data to develop a series of guiding questions that apply historical data to present fishery management: What were past conditions like under different climate

regimes? How did broad-scale climate changes affect biophysical, biological, and social dynamics in the marine environment? Can the answers to these questions provide a range of possible responses to changing climate conditions in increasingly warm periods, and be used to frame a range of possible responses?

These questions underscore that increasing ecological novelty does not necessarily mean a separation from historically continuous functions and composition (Hobbs et al. 2014, Heger et al. 2019). Even in cases where thresholds produce novel ecosystems or alternative stable states, there is much to be learned from historical data (Whitman et al. 2019). With a rise in unprecedented ecosystem conditions, historical knowledge can yield insights into the pathways of change or analogs offering clues for managing new conditions.

4. DISCUSSION AND CONCLUSIONS

During the past few decades, historical ecology has grown from an approach used by just a few scholars primarily studying terrestrial forests, to a large community of researchers and practitioners working around the world on a wide variety of terrestrial and aquatic organisms and ecosystems (Szabó & Hédl 2011, Rick & Lockwood 2013, Beller et al. 2020). Synthesis of publications focused on historical ecology demonstrates a dramatic increase in research during the past 20 yr (Fig. 1). A variety of initiatives, training opportunities, and professional networks (such as the Conservation Paleobiology Network and Oceans Past) will help carry forward historical ecology and its application to conservation. Given this increase in research and growing list of collaborative networks, our group of historical ecologists from around the world worked to assess the current state of historical ecology and provide future directions by developing research priorities centered around key issues and questions. Four key research priorities emerged: (1) methods and concepts, (2) knowledge co-production and community engagement, (3) policy and management, and (4) climate change, covering everything from machine learning and open access data, to diversifying perspectives, Indigenous knowledge, integrating disparate data sets and information, and the place of museums, social media, and other forms of engagement in broader education and research efforts. Here we discuss each of these priority areas and their value to historical ecology, conservation biology, and science more generally.

From history to paleoecology and archaeology, historical ecologists aim to understand natural climatic and anthropogenic environmental change, drawing on a wide range of different data sets and intellectual frameworks (Beller et al. 2020). Consequently, historical ecology is inherently interdisciplinary and requires collaboration that breaks down traditional intellectual silos. This is a key perspective that emerged in our 5 research questions focused on methods and concepts (Priority 1; see Box 1). Advancements in historical ecology demonstrate that both collaboration among scholars from diverse fields and the transdisciplinary expansion of individual scientists' interests and skills are important to effectively integrate quantitative and qualitative approaches (Allen 2007, Dietl & Flessa 2011, Clavero 2016, Clavero et al. 2016, Cramer et al. 2021, Dillon et al. 2022). While progress has been made, a need for better integration of data sets across spatio-temporal scales and drivers remains. At the same time, there is a call to move beyond data to take a critical view of baselines, considering the ways in which these are socially constructed, and the power dynamics embedded in the selection and implementation of baselines (Bliege Bird & Nimmo 2018, Soga & Gaston 2018, Collins et al. 2020, Duda et al. 2023, Palli et al. 2023). Historical ecology must continue to embrace its interdisciplinary nature and seek partnerships across disciplines, including natural science, social science, and the humanities, when developing and interrogating baselines and other aspects of research.

One of the most significant aspects of our research is the need for engaging diverse communities, the co-production of knowledge and research, and diversifying perspectives (Priority 2; see Box 1). This includes expanding training opportunities that promote diversity in scholarship and practice, enchainning funding opportunities for projects that emphasize under-represented groups and co-production, and democratizing knowledge through equitable open-access publication and dissemination. Our work here emphasizes the critical need for greater equity and engagement in historical ecology, which will ultimately enhance, expand, and improve research. While historical ecology is in many ways a leader of this type of research, there are still many areas for improvement, particularly to help make research less extractive and dominated by Western scientific interests and knowledge systems and towards one that appreciates diverse knowledge systems and is done with, for, and by Indigenous and other local communities (Liboiron 2021, de Vos & Schwartz 2022). Such a transdiscipli-

nary approach to historical ecology, based on knowledge co-production, fosters a holistic understanding of ecosystems and their dynamics, benefiting both scientific research and communities (Matarrita-Cascante et al. 2019, Dalton et al. 2020, Sidik 2022), and should be broadly incorporated in professional training, funding, and publication priorities. Continued recognition that people are not separate from ecosystems, not all human activities are negative, and people can be nested within ecosystems to promote long-term sustainability is important for better integrating distinct perspectives and knowledge systems to educate conservation practitioners, researchers, and the broader public. Knowledge co-production also seeks to break down traditional intellectual silos—a key goal of historical ecology—and one with significant opportunities for training, mentoring, and education. Finally, a more inclusive and equitable research framework offers a bridge between environmental justice, restoration, and ecological conservation with historical ecology poised to lead in these efforts (Douglass & Cooper 2020).

Historical ecology research priorities identified here also include the need for collaboration between academic researchers and practitioners (Priority 3). Curating data from the past that are relevant to conservation today demands collaboration with those implementing actions in relevant, applied management frameworks (Groff et al. 2023). Collaboration among researchers and practitioners overlaps with the previous research priorities by demonstrating another step in taking historical ecology from an academic pursuit to one focused on action and application to solving conservation and other environmental problems (Cavaleri Gerhardinger et al. 2023). Four of the questions identified in this priority emphasize the value of historical context as an integral part of the management decision-making process. Still, this is an area in urgent need of attention, especially investigations into how resource managers view findings from historical ecology and their application to decisions. Conservation efforts and application of historical ecological insights also draws from integrating diverse perspectives from different scientific communities and Indigenous communities into management decisions and priorities (Balée & Erickson 2006, Witter & Satterfield 2019).

Climate change is a central environmental concern of the 21st century, evidenced by 2023 being the hottest year on record, with a wide variety of climate-related perturbations, including massive fires in Canada and the Pacific, and extreme heat in American deserts and elsewhere, and marine heatwaves

globally (Asner et al. 2022, Speare et al. 2022, Tanaka & Van Houtan 2022). Priority 4 (see Box 1) emphasizes the contribution of historical ecology to climate change, discussed in 4 questions. One of the opportunities and challenges in this area is integrating long-term records of climate change that are generally on a global scale, with historical ecological data that are often more locally focused (Lotze et al. 2022, Palli et al. 2023). Similarly, the integration of distinct data sets focused on past climate such as fossil and marine sediment records provide opportunities to compare past climatic change and ecological responses, both with and without people, to help forecast future change (Harnik et al. 2012, Finnegan et al. 2024). Although we increasingly live in a no-analog world, historical ecological records still offer an unparalleled source of information on the relationships between climate change, anthropogenic processes, and the responses of ecosystems and organisms (Higgs et al. 2014).

The 4 research priorities discussed here demonstrate tremendous opportunity and growth for historical ecology. These priorities are all synergistic, illustrating the need to increase collaboration and expand historical ecology's field of inquiry, community of scholars, and practice and heighten inclusion and co-production of knowledge. These expanding frameworks will undoubtedly drive novel insights and breakthroughs and enhance the application of historical perspectives to contemporary and environmental issues, all while emphasizing the links between social justice and environmental conservation. Even though we are living on a rapidly changing planet, we believe that now more than ever, historical perspectives are central to helping better prepare for and navigate environmental uncertainty.

As we conducted our research, a commission of scholars was poised to mark the onset of a new geological epoch, the Anthropocene, or Age of Humans, that recognizes the profound influence of people on our planet (Lewis & Maslin 2015). After considerable debate, the committee rejected the proposal to set the beginning of the Anthropocene around 75 yr ago (~CE 1950), citing the longer time frame of human impacts on planetary processes, traced to the Industrial Revolution, colonizing the Americas and Australia, or onset of agriculture (Witze 2024). Historical perspectives from the preceding 12000 yr of the Holocene and earlier demonstrate that people influenced our planet for much longer than 75 yr, making perspectives from historical ecology crucial for understanding both how we arrived at present day conditions and illuminating the path ahead.

Data and material availability. All data needed to evaluate the conclusions in the paper are present in either the paper, the Supplementary Materials, or the linked repositories. Data and source code used in this study are available in the open-access third-party repository at GitHub (<https://bit.ly/477TePD>).

Acknowledgements. B. Godley, S. Pimm, and 2 anonymous reviewers improved earlier versions of this paper. We thank our home institutions, our collaborators, and a variety of funding agencies for supporting our fundamental research. This work was supported by the National Socio-Environmental Synthesis Center (SESYNC) under funding received from the National Science Foundation DBI-1639145. L.M. was supported by the Canada Research Chairs Program (CRC-2020-00204). R.H.T. was supported by the European Research Council (ERC) under the European Union's Horizon 2020 Research and Innovation Programme (# 856488). P.S. was supported by the Czech Academy of Sciences long-term research development project # RVO-67985939. A.C.C. was supported by the European Research Council (ERC) # 817911.

LITERATURE CITED

- ✦ Abu R, Reed MG, Jardine TD (2020) Using two-eyed seeing to bridge Western science and Indigenous knowledge systems and understand long-term change in the Saskatchewan River Delta, Canada. *Int J Water Resour Dev* 36:757–776
- ✦ Al-Abdulrazzak D, Pauly D (2017) Reconstructing historical baselines for the Persian/Arabian Gulf dugong, *Dugong dugon* (Mammalia: Sirena). *Zool Middle East* 63:95–102
- ✦ Alagona PS, Sandlos J, Wiersma YF (2012) Past imperfect: using historical ecology and baseline data for conservation and restoration projects in North America. *Environ Philos* 9:49–70
- ✦ Alagona P, Carey M, Howkins A (2023) Better together? The values, obstacles, opportunities, and prospects for collaborative research in environmental history. *Environ Hist* 28:269–299
- ✦ Allen MS (2007) Three millennia of human and sea turtle interactions in Remote Oceania. *Coral Reefs* 26:959–970
- ✦ Armstrong CG, Shoemaker AC, McKechnie I, Ekblom A and others (2017) Anthropological contributions to historical ecology: 50 questions, infinite prospects. *PLOS ONE* 12: e0171883
- ✦ Asner GP, Vaughn NR, Martin RE, Foo SA, Heckler J, Neilson BJ, Gove JM (2022) Mapped coral mortality and refugia in an archipelago-scale marine heat wave. *Proc Natl Acad Sci USA* 119:e2123331119
- ✦ Astudillo F, Becerra E, Delgado F, Jamieson R, Stahl PW (2023) When the archaeologists leave: legacies and services of the Historical Ecology of the Galápagos Islands Project. *Adv Archaeol Pract* 11:341–351
- ✦ Atalay S (2020) Indigenous science for a world in crisis. *Public Archaeol* 19:37–52
- ✦ Avtzi D, Stara K, Sgardeli V, Betsis A and others (2018) Quantifying the conservation value of Sacred Natural Sites. *Biol Conserv* 222:95–103
- ✦ Balée W (2006) The research program of historical ecology. *Annu Rev Anthropol* 35:75–98
- ✦ Balée W (2013) Cultural forests of the Amazon: a historical ecology of people and their landscapes. University of Alabama Press, Tuscaloosa, AL

- Balée W, Erickson CL (2006) Time, complexity, and historical ecology. In: Balée W, Erickson CL (eds) Time and complexity in historical ecology: studies in the Neotropical lowlands. Columbia University Press, New York, NY, p 1–20
- Barbeaux S, Aydin K, Fissel B, Holsman K and others (2019) Assessment of the Pacific cod stock in the Gulf of Alaska. North Pacific Fishery Management Council Gulf of Alaska Stock Assessment and Fishery Evaluation Report. <https://www.fisheries.noaa.gov/resource/data/2018-assessment-pacific-cod-stock-gulf-alaska>
- Barbeaux SJ, Holsman K, Zador S (2020) Marine heatwave stress test of ecosystem-based fisheries management in the Gulf of Alaska Pacific cod fishery. *Front Mar Sci* 7:703
- Barnosky AD, Hadly EA, Gonzalez P, Head J and others (2017) Merging paleobiology with conservation biology to guide the future of terrestrial ecosystems. *Science* 355: eaah4787
- Bartlett C, Marshall M, Marshall A (2012) Two-eyed seeing and other lessons learned within a co-learning journey of bringing together indigenous and mainstream knowledges and ways of knowing. *J Environ Stud Sci* 2: 331–340
- Beck J, Böller M, Erhardt A, Schwanghart W (2014) Spatial bias in the GBIF database and its effect on modeling species' geographic distributions. *Ecol Inform* 19:10–15
- Beller EE, McClenachan L, Zavaleta ES, Larsen LG (2020) Past forward: recommendations from historical ecology for ecosystem management. *Glob Ecol Conserv* 21: e00836
- Biernacki P, Waldorf D (1981) Snowball sampling: problems and techniques of chain referral sampling. *Sociol Methods Res* 10:141–163
- Bishop JC, Atkinson D, Walser RY (2013) The James Madison Carpenter Collection of Traditional Song and Drama. *Oral Tradit* 28:307–316
- Bliege Bird R, Bird DW (2021) Climate, landscape diversity, and food sovereignty in arid Australia: the firestick farming hypothesis. *Am J Hum Biol* 33:e23527
- Bliege Bird R, Nimmo D (2018) Restore the lost ecological functions of people. *Nat Ecol Evol* 2:1050–1052
- Bliege Bird R, Bird DW, Codding BF, Parker CH, Jones JH (2008) The 'fire stick farming' hypothesis: Australian Aboriginal foraging strategies, biodiversity, and anthropogenic fire mosaics. *Proc Natl Acad Sci USA* 105: 14796–14801
- Bliege Bird R, Taylor N, Codding BF, Bird DW (2013) Niche construction and Dreaming logic: aboriginal patch mosaic burning and varanid lizards (*Varanus gouldii*) in Australia. *Proc R Soc B* 280:20132297
- Bliege Bird R, Bird DW, Codding BF (2016) People, El Niño southern oscillation and fire in Australia: fire regimes and climate controls in hummock grasslands. *Philos Trans R Soc B* 371:20150343
- Bluhm LE, Surovell TA (2019) Validation of a global model of taphonomic bias using geologic radiocarbon ages. *Quat Res* 91:325–328
- Boivin NL, Zeder MA, Fuller DQ, Crowther A and others (2016) Ecological consequences of human niche construction: examining long-term anthropogenic shaping of global species distributions. *Proc Natl Acad Sci USA* 113:6388–6396
- Boles OJ, Shoemaker A, Courtney Mustaphi CJ, Petek N, Ekblom A, Lane PJ (2019) Historical ecologies of pastoralist overgrazing in Kenya: long-term perspectives on cause and effect. *Hum Ecol* 47:419–434
- Braje TJ, Rick TC, Erlandson JM, Rogers-Bennett L, Catton CA (2016) Historical ecology can inform restoration site selection: the case of black abalone (*Haliotis cracherodii*) along California's Channel Islands. *Aquat Conserv* 26: 470–481
- Buck M, Hamilton C (2011) The Nagoya Protocol on access to genetic resources and the fair and equitable sharing of benefits arising from their utilization to the Convention on Biological Diversity. *Rev Eur Community Int Environ Law* 20:47–61
- Burns F, August T, Eaton M, Noble D, Powney G, Isaac N, Hayhow D (2023) UK biodiversity indicators. Joint Nature Conservation Committee, Peterborough
- Carmenta R, Vira B (2018) Integration for restoration: reflecting on lessons learned from the silos of the past. In: Mansourian S, Parrotta J (eds) Forest landscape restoration: integrated approaches to support effective implementation. Routledge, London, p 16–36
- Carroll SR, Garba I, Figueroa-Rodriguez OL, Holbrook J, and others (2020) The CARE principles for Indigenous data governance. *Data Sci J* 19:43
- Cavaleri Gerhardinger L, Brodie Rudolph T, Gaill F, Mortyn G and others (2023) Bridging shades of blue: co-constructing knowledge with the International Panel for Ocean Sustainability. *Coast Manage* 51:244–264
- Christenson AJ (2007) *Popol Vuh: the sacred book of the Maya, Vol 1*. University of Oklahoma Press, Norman, OK
- Clark TR, Roff G, Zhao J, Feng Y, Done TJ, Pandolfi JM (2014) Testing the precision and accuracy of the U–Th chronometer for dating coral mortality events in the last 100 years. *Quat Geochronol* 23:35–45
- Clavero M (2016) Species substitutions driven by anthropogenic positive feedbacks: Spanish crayfish species as a case study. *Biol Conserv* 193:80–85
- Clavero M, Nores C, Kubersky-Piredda S, Centeno-Cuadros A (2016) Interdisciplinarity to reconstruct historical introductions: solving the status of cryptogenic crayfish. *Biol Rev Camb Philos Soc* 91:1036–1049
- Clement CR, Denevan WM, Heckenberger MJ, Junqueira AB, Neves EG, Teixeira WG, Woods WI (2015) The domestication of Amazonia before European conquest. *Proc R Soc B* 282:20150813
- Codding BF, Bliege Bird R, Kauhanen PG, Bird DW (2014) Conservation or co-evolution? Intermediate levels of aboriginal burning and hunting have positive effects on kangaroo populations in Western Australia. *Hum Ecol* 42:659–669
- Coleman MA, Wood G, Filbee-Dexter K, Minne AJP and others (2020) Restore or redefine: future trajectories for restoration. *Front Mar Sci* 7:237
- Collins AC, Böhm M, Collen B (2020) Choice of baseline affects historical population trends in hunted mammals of North America. *Biol Conserv* 242:108421
- Colonese AC, Brugere C, Ramires M, Clauzet M and others (2023) The legacy of pre-Columbian fisheries to food security and poverty alleviation in the modern Amazon. In: Colonese AC, Milheira RG (eds) Historical ecology and landscape archaeology in Lowland South America. Springer International Publishing, Cham, p 3–19
- Cox KD, Gerwing TG, Macdonald T, Hessing-Lewis M and others (2019) Infaunal community responses to ancient clam gardens. *ICES J Mar Sci* 76:2362–2373
- Cramer KL, Donovan MK, Jackson JBC, Greenstein BJ, Korpanty CA, Cook GM, Pandolfi JM (2021) The transformation of Caribbean coral communities since humans. *Ecol Evol* 11:10098–10118

- Crowley TJ, Berner RA (2001) CO₂ and climate change. *Science* 292:870–872
- Crumley CL (ed) (1994) *Historical ecology: cultural knowledge and changing landscapes*. School of American Research Press, Santa Fe, NM
- Crumley CL (2021) Historical ecology: a robust bridge between archaeology and ecology. *Sustainability* 13:8210
- Crystal-Ornelas R, Varadharajan C, O’Ryan D, Beilsmith K and others (2022) Enabling FAIR data in Earth and environmental science with community-centric (meta)data reporting formats. *Sci Data* 9:700
- Curran K, Zimmermann N (2022) The dynamics of collaboration in heritage science. *Stud Conserv* 67:136–149
- Dalton K, Skrobe M, Bell H, Kantner B, Berndtson D, Gerhardinger LC, Christie P (2020) Marine-related learning networks: shifting the paradigm toward collaborative ocean governance. *Front Mar Sci* 7:595054
- Daru BH, Rodriguez J (2023) Mass production of unvouchered records fails to represent global biodiversity patterns. *Nat Ecol Evol* 7:816–831
- Davidson E, Edwards R, Jamieson L, Weller S (2019) Big data, qualitative style: a breadth-and-depth method for working with large amounts of secondary qualitative data. *Qual Quant* 53:363–376
- Davis AO (2018) *Changing perspectives on citizen science using eBird data on Grand Bahama Island, The Bahamas*. PhD dissertation, Miami University, Oxford, OH
- Davison CW, Rahbek C, Morueta-Holme N (2021) Land-use change and biodiversity: challenges for assembling evidence on the greatest threat to nature. *Glob Change Biol* 27:5414–5429
- Dawson NM, Coolsaet B, Sterling EJ, Loveridge R and others (2021) The role of Indigenous peoples and local communities in effective and equitable conservation. *Ecol Soc* 26:19
- de Vos A, Schwartz MW (2022) Confronting parachute science in conservation. *Conserv Sci Pract* 4:e12681
- Degroot D, Anchukaitis K, Bauch M, Burnham J and others (2021) Towards a rigorous understanding of societal responses to climate change. *Nature* 591:539–550
- Denevan WM (1992) The pristine myth: the landscape of the Americas in 1492. *Ann Assoc Am Geogr* 82:369–385
- DeSilvey C, Harrison R (2020) Anticipating loss: rethinking endangerment in heritage futures. *Int J Heritage Stud* 26: 1–7
- Desse-Berset N (2011) Ancient sturgeon populations in France through archaeozoological remains, from prehistoric time until the eighteenth century. In: Williot P, Rochard E, Desse-Berset N, Kirschbaum F, Gessner J (eds) *Biology and conservation of the European sturgeon *Acipenser sturio* L. 1758: the reunion of the European and Atlantic sturgeons*. Springer, Berlin, p 91–115
- Deutsch C, Ferrel A, Seibel B, Pörtner HO, Huey RB (2015) Climate change tightens a metabolic constraint on marine habitats. *Science* 348:1132–1135
- Dieltl GP, Flessa KW (2011) Conservation paleobiology: putting the dead to work. *Trends Ecol Evol* 26:30–37
- Dillon EM, Lafferty KD, McCauley DJ, Bradley D and others (2020) Dermal denticle assemblages in coral reef sediments correlate with conventional shark surveys. *Methods Ecol Evol* 11:362–375
- Dillon EM, Pier JQ, Smith JA, Raja NB and others (2022) What is conservation paleobiology? Tracking 20 years of research and development. *Front Ecol Evol* 10:1031483
- Dorfman E (2017) *The future of natural history museums*. Routledge, London
- Douglass K, Cooper J (2020) Archaeology, environmental justice, and climate change on islands of the Caribbean and southwestern Indian Ocean. *Proc Natl Acad Sci USA* 117:8254–8262
- Drotner K, Dziekan V, Parry R, Schrøder KC (2019) *The Routledge handbook of museums, media and communication*. Routledge, London
- Duda MP, Grooms C, Sympson L, Blais JM and others (2023) A 2200-year record of Andean Condor diet and nest site usage reflects natural and anthropogenic stressors. *Proc R Soc B* 290:20230106
- Early-Capistrán MM, Sáenz-Arroyo A, Cardoso-Mohedano JG, Garibay-Melo G, Peckham SH, Koch V (2018) Reconstructing 290 years of a data-poor fishery through ethnographic and archival research: the East Pacific green turtle (*Chelonia mydas*) in Baja California, Mexico. *Fish Fish* 19:57–77
- Early-Capistrán MM, Solana-Arellano E, Abreu-Grobois FA, Narchi NE and others (2020) Quantifying local ecological knowledge to model historical abundance of long-lived, heavily-exploited fauna. *PeerJ* 8:e9494
- Ellis E, Maslin M, Boivin N, Bauer A (2016) Involve social scientists in defining the Anthropocene. *Nature* 540: 192–193
- Ellis EC, Gauthier N, Klein Goldewijk K, Bliege Bird R and others (2021) People have shaped most of terrestrial nature for at least 12,000 years. *Proc Natl Acad Sci USA* 118:e2023483118
- Emslie SD (2021) Ancient Adélie penguin colony revealed by snowmelt at Cape Irizar, Ross Sea, Antarctica. *Geology* 49:145–149
- Evans SM, Gebbels S, Stockill JM (2008) ‘Our shared responsibility’: participation in ecological projects as a means of empowering communities to contribute to coastal management processes. *Mar Pollut Bull* 57:3–7
- Evans HJ, Nicolaisen L, Tougaard S, Achiam M (2020) Perspective. Museums beyond neutrality. *Nord Museol* 2: 19–25
- Felton A, Fischer J, Lindenmayer DB, Montague-Drake R and others (2009) Climate change, conservation and management: an assessment of the peer-reviewed scientific journal literature. *Biodivers Conserv* 18:2243–2253
- Fernando T (2022) Seeing like the sea: a multispecies history of the Ceylon pearl fishery 1800–1925. *Past Present* 254: 127–160
- Finnegan S, Anderson SC, Harnik PG, Simpson C and others (2015) Paleontological baselines for evaluating extinction risk in the modern oceans. *Science* 348:567–570
- Finnegan S, Harnik PG, Lockwood R, Lotze HK, McClenahan L, Kahanamoku SS (2024) Using the fossil record to understand extinction risk and inform climate-adapted conservation in a changing ocean. *Annu Rev Mar Sci* 16: 307–333
- Fischer J, Riechers M, Loos J, Martin-Lopez B, Tempterton VM (2021) Making the UN Decade on Ecosystem Restoration a social-ecological endeavour. *Trends Ecol Evol* 36: 20–28
- Fisk JJ, Jacobs LA, Russo BUK, Meier E and others (2021) Cultivating sovereignty in parks and protected areas: sowing the seeds of restorative and transformative justice through the #LANDBACK movement. *Parks Stewardship Forum* 37:517–526
- Flantua SGA, O’Dea A, Onstein RE, Giraldo C, Hooghiemstra H (2019) The flickering connectivity system of the north Andean páramos. *J Biogeogr* 46:1808–1825
- Fletcher MS, Hamilton R, Dressler W, Palmer L (2021)

- Indigenous knowledge and the shackles of wilderness. *Proc Natl Acad Sci USA* 118:e2022218118
- ✦ Forcina G, Guerrini M, van Grouw H, Gupta BK and others (2015) Impacts of biological globalization in the Mediterranean: unveiling the deep history of human-mediated gamebird dispersal. *Proc Natl Acad Sci USA* 112: 3296–3301
- ✦ Fossile T, Herbst DF, McGrath K, Toso A and others (2023) Bridging archaeology and marine conservation in the Neotropics. *PLOS ONE* 18:e0285951
- ✦ Fuchs R, Herold M, Verburg PH, Clevers JGPW, Eberle J (2015) Gross changes in reconstructions of historic land cover/use for Europe between 1900 and 2010. *Glob Change Biol* 21:299–313
- ✦ Gagné TO, Reygondeau G, Jenkins CN, Sexton JO, Bograd SJ, Hazen EL, Van Houtan KS (2020) Towards a global understanding of the drivers of marine and terrestrial biodiversity. *PLOS ONE* 15:e0228065
- Galla CK (2009) Indigenous language revitalization and technology from traditional to contemporary domains. In: Reyhner J, Lockard L (eds) *Indigenous language revitalization: encouragement, guidance & lessons learned*. Northern Arizona University, Flagstaff, AZ, p 167–182
- ✦ Garnett ST, Burgess ND, Fa JE, Fernández-Llamazares Á and others (2018) A spatial overview of the global importance of Indigenous lands for conservation. *Nat Sustain* 1: 369–374
- ✦ Gerhardinger LC, de Andrade MM, Corrêa MR, Turra A (2020) Crafting a sustainability transition experiment for the Brazilian blue economy. *Mar Policy* 120:104157
- Gessner J, Arndt GM, Fredrich F, Ludwig A, Kirschbaum F, Bartel R, Nordheim H (2011) Remediation of Atlantic sturgeon *Acipenser oxyrinchus* in the Oder River: background and first results. In: Williot P, Rochard E, Dese-Berset N, Kirschbaum F, Gessner J (eds) *Biology and conservation of the European Sturgeon *Acipenser sturio* L. 1758: the reunion of the European and Atlantic sturgeons*. Springer, Berlin, p 539–559
- ✦ Gil-Romera G, Lamb HF, Turton D, Sevilla-Callejo M, Umer M (2010) Long-term resilience, bush encroachment patterns and local knowledge in a Northeast African savanna. *Glob Environ Change* 20:612–626
- ✦ Gil-Romera G, Turton D, Sevilla-Callejo M (2011) Landscape change in the lower Omo valley, southwestern Ethiopia: burning patterns and woody encroachment in the savanna. *J East Afr Stud* 5:108–128
- ✦ Gillies CL, Castine SA, Alleyway HK, Crawford C and others (2020) Conservation status of the oyster reef ecosystem of southern and eastern Australia. *Glob Ecol Conserv* 22: e00988
- ✦ Grace MK, Akçakaya HR, Bennett EL, Brooks TM and others (2021) Testing a global standard for quantifying species recovery and assessing conservation impact. *Conserv Biol* 35:1833–1849
- ✦ Gray RRR (2022) Rematriation: Ts'msyen law, rights of relationality, and protocols of return. *Native Am Indig Stud* 9: 1–27
- ✦ Grayson DK (2005) A brief history of Great Basin pikas. *J Biogeogr* 32:2103–2111
- ✦ Greer S (2010) Heritage and empowerment: community-based Indigenous cultural heritage in northern Australia. *Int J Herit Stud* 16:45–58
- ✦ Groff DV, McDonough MacKenzie C, Pier JQ, Shaffer AB, Dietl GP (2023) Knowing but not doing: quantifying the research-implementation gap in conservation paleobiology. *Front Ecol Evol* 11:1058992
- ✦ Hall L, Dell CA, Fornssler B, Hopkins C, Mushquash C, Rowan M (2015) Research as cultural renewal: applying two-eyed seeing in a research project about cultural interventions in First Nations addictions treatment. *Int Indig Policy J* 6:1–15
- ✦ Harnik PG, Lotze HK, Anderson SC, Finkel ZV and others (2012) Extinctions in ancient and modern seas. *Trends Ecol Evol* 27:608–617
- ✦ Harris LA, Garza C, Hatch M, Parrish J and others (2021) Equitable exchange: a framework for diversity and inclusion in the geosciences. *AGU Advances* 2:e2020 AV000359
- Hassan F (2014) Tangible heritage in archaeology. In: Smith C (ed) *Encyclopedia of global archaeology*. Springer, New York, NY, p 10489–10492
- Hassan F (2021) Cultural heritage, empowerment and the social transformation of local communities. In: Higgins V, Douglas D (eds) *Communities and cultural heritage: global issues, local values*. Routledge, London, p 23–35
- ✦ Hatch MBA, Parrish JK, Heppell SS, Augustine S and others (2023) Boundary spanners: a critical role for enduring collaborations between Indigenous communities and mainstream scientists. *Ecol Soc* 28:41
- ✦ Hédl R, Kopecký M, Komárek J (2010) Half a century of succession in a temperate oakwood: from species-rich community to mesic forest. *Divers Distrib* 16:267–276
- ✦ Heger T, Bernard-Verdier M, Gessler A, Greenwood AD and others (2019) Towards an integrative, eco-evolutionary understanding of ecological novelty: studying and communicating interlinked effects of global change. *BioScience* 69:888–899
- ✦ Herbst DF, Rampon J, Baleeiro B, Silva LG, Fossile T, Colonese AC (2023) 180 years of marine animal diversity as perceived by public media in southern Brazil. *PLOS ONE* 18:e0284024
- ✦ Higgs E, Falk DA, Guerrini A, Hall M and others (2014) The changing role of history in restoration ecology. *Front Ecol Environ* 12:499–506
- Hilding-Rydevik T, Moen J, Green C (2017) Baselines and the shifting baseline syndrome — exploring frames of reference in nature conservation. In: Westin A, Crumley CL, Lennartsson T (eds) *Issues and concepts in historical ecology: the past and future of landscapes and regions*. Cambridge University Press, Cambridge, p 112–142
- ✦ Hill R, Cullen-Unsworth LC, Talbot LD, McIntyre-Tamwoy S (2011) Empowering Indigenous peoples' biocultural diversity through World Heritage cultural landscapes: a case study from the Australian humid tropical forests. *Int J Herit Stud* 17:571–591
- ✦ Hobbs RJ, Higgs E, Hall CM, Bridgewater P and others (2014) Managing the whole landscape: historical, hybrid, and novel ecosystems. *Front Ecol Environ* 12:557–564
- ✦ Hoffman KM, Davis EL, Wickham SB, Schang K and others (2021) Conservation of Earth's biodiversity is embedded in Indigenous fire stewardship. *Proc Natl Acad Sci USA* 118:e2105073118
- ✦ Hortal J, Jiménez-Valverde A, Gómez JF, Lobo JM, Baselga A (2008) Historical bias in biodiversity inventories affects the observed environmental niche of the species. *Oikos* 117:847–858
- ✦ Hughes AC, Orr MC, Ma K, Costello MJ and others (2021) Sampling biases shape our view of the natural world. *Ecography* 44:1259–1269
- ISE (2006) International Society of Ethnobiology code of ethics (with 2008 additions). International Society of Ethnobiology, Gainesville, FL

- IUCN Species Survival Commission (2021) IUCN green status of species. International Union for Conservation of Nature, Gland
- ✦ Jackson JBC (1997) Reefs since Columbus. *Coral Reefs* 16: S23–S32
- ✦ Jackson JBC, Erwin DH (2006) What can we learn about ecology and evolution from the fossil record? *Trends Ecol Evol* 21:322–328
- ✦ Jackson JBC, Kirby MX, Berger WH, Bjorndal KA and others (2001) Historical overfishing and the recent collapse of coastal ecosystems. *Science* 293:629–637
- ✦ Jackson ST, Sax DF (2010) Balancing biodiversity in a changing environment: extinction debt, immigration credit and species turnover. *Trends Ecol Evol* 25:153–160
- Janes RR, Sandell R (2019) *Museum activism*. Routledge, London
- ✦ Jerardino A (2012) What archaeology can tell us about sustainability and climate change: a South African west coast perspective. *J Mar Sci Res Dev* 1:105
- ✦ Jindal A (2022) Misguided artificial intelligence: how racial bias is built into clinical models. *J Brown Hosp Med* 2: 1–5
- ✦ Jorgensen SJ, Micheli F, White TD, Van Houtan KS and others (2022) Emergent research and priorities for shark and ray conservation. *Endang Species Res* 47:171–203
- ✦ Jung MR, Horgen FD, Orski SV, Rodriguez V and others (2018) Validation of ATR FT-IR to identify polymers of plastic marine debris, including those ingested by marine organisms. *Mar Pollut Bull* 127:704–716
- ✦ Kidwell SM (2015) Biology in the Anthropocene: challenges and insights from young fossil records. *Proc Natl Acad Sci USA* 112:4922–4929
- ✦ Kittinger JN, Van Houtan KS, McClenachan LE, Lawrence AL (2013) Using historical data to assess the biogeography of population recovery. *Ecography* 36:868–872
- Kittinger JN, McClenachan L, Gedan KB, Blight LK (2015) *Marine historical ecology in conservation: applying the past to manage for the future*. University of California Press, Oakland, CA
- ✦ Klenk N, Fiume A, Meehan K, Gibbes C (2017) Local knowledge in climate adaptation research: moving knowledge frameworks from extraction to co-production. *Wiley Interdiscip Rev Clim Change* 8:e475
- ✦ Koslov L (2016) The case for retreat. *Public Cult* 28:359–387
- ✦ Lawson KN, Letendre H, Drew JA (2021) Historical maps provide insight into a century and a half of habitat change in Fijian coasts. *Ecol Evol* 11:15573–15584
- Leonard K, David-Chavez D, Smiles D, Jennings L and others (2023) Water back: a review centering rematriation and indigenous water research sovereignty. *Water Altern* 16: 374–428
- Leopold AS, Cain SA, Cottam CM, Gabrielson IN, Kimball TL (1963) *Wildlife management in the national parks*. National Parks Service, Washington, DC
- ✦ Lepofsky D, Toniello G, Earnshaw J, Roberts C, Wilson L, Rowell K, Holmes K (2021) Ancient anthropogenic clam gardens of the northwest coast expand clam habitat. *Ecosystems* 24:248–260
- ✦ Levine J, Muthukrishna M, Chan KMA, Satterfield T (2017) Sea otters, social justice, and ecosystem-service perceptions in Clayoquot Sound, Canada. *Conserv Biol* 31: 343–352
- ✦ Levis C, Costa FRC, Bongers F, Peña-Claros M and others (2017) Persistent effects of pre-Columbian plant domestication on Amazonian forest composition. *Science* 355: 925–931
- ✦ Lewis SL, Maslin MA (2015) Defining the Anthropocene. *Nature* 519:171–180
- ✦ Lewison R, Oro D, Godley BJ, Underhill L and others (2012) Research priorities for seabirds: improving conservation and management in the 21st century. *Endang Species Res* 17:93–121
- Liboiron M (2021) *Pollution is colonialism*. Duke University Press, Durham, NC
- Linstone HA, Tuross M (1975) *The Delphi method*. Addison-Wesley, Reading, MA
- ✦ Lombardo U, Szabo K, Capriles JM, May JH and others (2013) Early and Middle Holocene hunter-gatherer occupations in western Amazonia: the hidden shell middens. *PLOS ONE* 8:e72746
- ✦ Lombardo U, Iriarte J, Hilbert L, Ruiz-Pérez J, Capriles JM, Veit H (2020) Early Holocene crop cultivation and landscape modification in Amazonia. *Nature* 581:190–193
- Lotze HK, McClenachan L (2013) *Marine historical ecology: informing the future by learning from the past*. In: Bertness MD, Bruno JF, Silliman BR, Stachowicz JJ (eds) *Marine community ecology and conservation*. Sinauer, Sunderland, MA, p 165–201
- ✦ Lotze HK, Lenihan HS, Bourque BJ, Bradbury RH and others (2006) Depletion, degradation, and recovery potential of estuaries and coastal seas. *Science* 312:1806–1809
- ✦ Lotze HK, Coll M, Dunne JA (2011a) Historical changes in marine resources, food-web structure and ecosystem functioning in the Adriatic Sea, Mediterranean. *Ecosystems* 14:198–222
- Lotze HK, Erlandson JM, Hardt MJ, Norris RD, Roy K, Smith TD, Whitcraft CR (2011b) *Uncovering the ocean's past*. In: Jackson JBC, Alexander KE, Sala E (eds) *Shifting baselines: the past and the future of ocean fisheries*. Island Press, Washington, DC, p 137–161
- ✦ Lotze HK, Tittensor DP, Bryndum-Buchholz A, Eddy TD and others (2019) Global ensemble projections reveal trophic amplification of ocean biomass declines with climate change. *Proc Natl Acad Sci USA* 116:12907–12912
- ✦ Lotze HK, Mellon S, Coyne J, Betts M and others (2022) Long-term ocean and resource dynamics in a hotspot of climate change. *Facets* 7:1142–1184
- ✦ Ludwig A, Debus L, Lieckfeldt D, Wirgin I and others (2002) When the American sea sturgeon swam east. *Nature* 419: 447–448
- ✦ Lydersen JM, Collins BM (2018) Change in vegetation patterns over a large forested landscape based on historical and contemporary aerial photography. *Ecosystems* 21: 1348–1363
- ✦ Lyman RL (2012) A warrant for applied palaeozoology. *Biol Rev Camb Philos Soc* 87:513–525
- ✦ Mace GM (2014) Whose conservation? *Science* 345: 1558–1560
- ✦ Malsale P, Sanau N, Tofaeono TI, Kavisi Z and others (2018) Protocols and partnerships for engaging Pacific Island communities in the collection and use of traditional climate knowledge. *Bull Am Meteorol Soc* 99:2471–2489
- Marshall M, Marshall A, Bartlett C (2015) *Two-eyed seeing in medicine*. In: Greenwood M, de Leeuw S, Lindsay NM (eds) *Determinants of Indigenous peoples' health in Canada: beyond the social*, 2nd edn. Canadian Scholar's Press, Toronto, p 16–24
- ✦ Martín-Martín A, Orduna-Malea E, Thelwall M, Delgado López-Cózar E (2018) Google Scholar, Web of Science, and Scopus: a systematic comparison of citations in 252 subject categories. *J Informetrics* 12:1160–1177
- ✦ Matarrita-Cascante D, Sene-Harper A, Ruyle L (2019) A

- holistic framework for participatory conservation approaches. *Int J Sustain Dev World Ecol* 26:484–494
- ✦ McAfee D, Reinhold SL, Alleway HK, Connell SD (2021) Environmental solutions fast-tracked: reversing public scepticism to public engagement. *Biol Conserv* 253: 108899
- ✦ McAfee D, McLeod IM, Alleway HK, Bishop MJ and others (2022) Turning a lost reef ecosystem into a national restoration program. *Conserv Biol* 36:e13958
- ✦ McClenachan L (2009a) Documenting loss of large trophy fish from the Florida Keys with historical photographs. *Conserv Biol* 23:636–643
- ✦ McClenachan L (2009b) Historical declines of goliath grouper populations in South Florida, USA. *Endang Species Res* 7:175–181
- ✦ McClenachan L, Cooper AB (2008) Extinction rate, historical population structure and ecological role of the Caribbean monk seal. *Proc R Soc B* 275:1351–1358
- ✦ McClenachan L, Ferretti F, Baum JK (2012) From archives to conservation: why historical data are needed to set baselines for marine animals and ecosystems. *Conserv Lett* 5: 349–359
- ✦ McClenachan L, Cooper AB, McKenzie MG, Drew JA (2015) The importance of surprising results and best practices in historical ecology. *BioScience* 65:932–939
- ✦ McKechnie I, Lepofsky D, Moss ML, Butler VL and others (2014) Archaeological data provide alternative hypotheses on Pacific herring (*Clupea pallasii*) distribution, abundance, and variability. *Proc Natl Acad Sci USA* 111: E807–E816
- McKeon M (2012) Two-eyed seeing into environmental education: revealing its “natural” readiness to indigenize. *Can J Environ Educ* 17:131–147
- ✦ McNiven IJ, Bedingfield AC (2008) Past and present marine mammal hunting rates and abundances: dugong (*Dugong dugon*) evidence from Dabangai Bone Mound, Torres Strait. *J Archaeol Sci* 35:505–515
- ✦ Meagher K (2021) Introduction: the politics of open access—decolonizing research or corporate capture? *Dev Change* 52:340–358
- ✦ Meltzer DJ (1999) Human responses to middle Holocene (Altitheal) climates on the north American Great Plains. *Quat Res* 52:404–416
- ✦ Millar CI, Stephenson NL, Stephens SL (2007) Climate change and forests of the future: managing in the face of uncertainty. *Ecol Appl* 17:2145–2151
- ✦ Miller EA, McClenachan L, Uni Y, Phocas G, Hagemann ME, Van Houtan KS (2019) The historical development of complex global trafficking networks for marine wildlife. *Sci Adv* 5:eaav5948
- ✦ Miller EA, Lisin SE, Smith CM, Van Houtan KS (2020) Herbaria macroalgae as a proxy for historical upwelling trends in Central California. *Proc R Soc B* 287:20200732
- ✦ Miller EA, Yamahara KM, French C, Spingarn N, Birch JM, Van Houtan KS (2022) A Raman spectral reference library of potential anthropogenic and biological ocean polymers. *Sci Data* 9:780
- ✦ Mistry J, Berardi A (2016) Bridging indigenous and scientific knowledge. *Science* 352:1274–1275
- ✦ Morgan RM, Kneebone RL, Pyenson ND, Sholts SB, Houstoun W, Butler B, Chesters K (2023) Regaining creativity in science: insights from conversation. *R Soc Open Sci* 10: 230134
- ✦ Moutinho S (2022) Brazil's election is a cliffhanger for scientists. *Science* 378:235–236
- ✦ Munteanu C, Kuemmerle T, Keuler NS, Müller D and others (2015) Legacies of 19th century land use shape contemporary forest cover. *Glob Environ Change* 34:83–94
- ✦ Munteanu C, Kamp J, Nita MD, Klein N and others (2020) Cold War spy satellite images reveal long-term declines of a philopatric keystone species in response to cropland expansion. *Proc R Soc B* 287:20192897
- ✦ Mychajliw AM, Ellwood ER, Alagona PS, Anderson RS and others (2022) Lessons for conservation from beneath the pavement. *Conserv Biol* 36:e13983
- Nicholson TE, McClenachan L, Tanaka KR, Van Houtan KS (2024) Sea otter recovery buffers century-scale declines in California kelp forests. *PLOS Clim* 3:e0000290
- NOAA (2020) NOAA data strategy: maximizing the value of NOAA data. NOAA Science & Technology Focus Areas. National Oceanic and Atmospheric Administration, Silver Spring, MD
- ✦ Nolan C, Overpeck JT, Allen JR, Anderson PM and others (2018) Past and future global transformation of terrestrial ecosystems under climate change. *Science* 361:920–923
- ✦ Obermeyer Z, Powers B, Vogeli C, Mullainathan S (2019) Dissecting racial bias in an algorithm used to manage the health of populations. *Science* 366:447–453
- ✦ Palli J, Mensing SA, Schoolman EM, Solano F, Piovesan G (2023) Historical ecology identifies long-term rewilding strategy for conserving Mediterranean mountain forests in south Italy. *Ecol Appl* 33:e2758
- ✦ Palomares MLD, Heymans JJ, Pauly D (2007) Historical ecology of the Raja Ampat Archipelago, Papua Province, Indonesia. *Hist Philos Life Sci* 29:33–56
- ✦ Pandolfi JM, Bradbury RH, Sala E, Hughes TP and others (2003) Global trajectories of the long-term decline of coral reef ecosystems. *Science* 301:955–958
- ✦ Paulson T, Brown KC, Alagona PS (2021) The test of time: using historical methods to assess models of ecological change on California's hardwood rangelands. *Ann Assoc Am Geogr* 111:402–421
- ✦ Pauly D (1995) Anecdotes and the shifting baseline syndrome of fisheries. *Trends Ecol Evol* 10:430
- ✦ Peacock E, Randklev CR, Wolverton S, Palmer RA, Zaleski S (2012) The ‘cultural filter,’ human transport of mussel shell, and the applied potential of zooarchaeological data. *Ecol Appl* 22:1446–1459
- ✦ Piñar G, Sterflinger K (2021) Natural sciences at the service of art and cultural heritage: an interdisciplinary area in development and important challenges. *Microb Biotechnol* 14:806–809
- ✦ Polónia A (2020) Inter-, multi- and trans-disciplinarity in maritime history: potentialities and limits. *Int J Marit Hist* 32: 414–425
- ✦ Pooley S (2013) Historians are from Venus, ecologists are from Mars. *Conserv Biol* 27:1481–1483
- R Core Team (2023) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna
- ✦ Raja NB, Dunne EM, Matiwane A, Khan TM, Nätscher PS, Ghilardi AM, Chattopadhyay D (2022) Colonial history and global economics distort our understanding of deep-time biodiversity. *Nat Ecol Evol* 6:145–154
- ✦ Ratnasingham S, Hebert PDN (2007) BOLD: the Barcode of Life data system. *Mol Ecol Notes* 7:355–364
- ✦ Reddin CJ, Aberhan M, Raja NB, Kocsis ÁT (2022) Global warming generates predictable extinctions of warm- and cold-water marine benthic invertebrates via thermal habitat loss. *Glob Change Biol* 28:5793–5807
- ✦ Reeder-Myers L, Braje TJ, Hofman CA, Elliott Smith EA and others (2022) Indigenous oyster fisheries persisted for

- millennia and should inform future management. *Nat Commun* 13:2383
- Rees AF, Alfaro-Shigueto J, Barata PCR, Bjorndal KA and others (2016) Are we working towards global research priorities for management and conservation of sea turtles? *Endang Species Res* 31:337–382
- Reid DA, Vail DD (2019) Interpreting the environment at museums and historic sites. *Interpreting History Series*, Vol 17. Rowman & Littlefield, Lanham, MD
- Reid AJ, Eckert LE, Lane JF, Young N and others (2021) 'Two-eyed seeing': an indigenous framework to transform fisheries research and management. *Fish Fish* 22: 243–261
- Reif J, Szarvas F, Šťastný K (2021) 'Tell me where the birds have gone'—reconstructing historical influence of major environmental drivers on bird populations from memories of ornithologists of an older generation. *Ecol Indic* 129:107909
- Reyes-García V (2024) *Routledge handbook of climate change impacts on indigenous peoples and local communities*. Routledge, New York, NY
- Reygondeau G, Cheung WWL, Wabnitz CCC, Lam VWY, Frölicher T, Maury O (2020) Climate change-induced emergence of novel biogeochemical provinces. *Front Mar Sci* 7:657
- Riccaboni M, Verginer L (2022) The impact of the COVID-19 pandemic on scientific research in the life sciences. *PLOS ONE* 17:e0263001
- Rick TC, Lockwood R (2013) Integrating paleobiology, archeology, and history to inform biological conservation. *Conserv Biol* 27:45–54
- Rick T, Ontiveros MÁC, Jerardino A, Mariotti A, Méndez C, Williams AN (2020) Human-environmental interactions in Mediterranean climate regions from the Pleistocene to the Anthropocene. *Anthropocene* 31:100253
- Rivera-Collazo IC, Perdikaris S (2023) Climate change, site formation, and indigenous use of coastlines in Barbuda. *Holocene* 33:1142–1153
- Rizayeva A, Nita MD, Radloff VC (2023) Large-area, 1964 land cover classifications of Corona spy satellite imagery for the Caucasus Mountains. *Remote Sens Environ* 284: 113343
- Robinson H (2017) Is cultural democracy possible in a museum? Critical reflections on Indigenous engagement in the development of the exhibition *Encounters: revealing Stories of Aboriginal and Torres Strait Islander Objects from the British Museum*. *Int J Herit Stud* 23:860–874
- Rose DB (2004) *Reports from a wild country: ethics for decolonisation*. UNSW Press, Sydney
- Rost D (2018) *Shifting baselines: interdisciplinary perspectives on long-term change perception and memory*. Ostfalia University of Applied Sciences, Wolfenbüttel
- Sáenz-Arroyo A, Revollo-Fernández D (2016) Local ecological knowledge concurs with fishing statistics: an example from the abalone fishery in Baja California, Mexico. *Mar Policy* 71:217–221
- Salomon AK, Wilson KBJ, White XE, Tanape N Sr, Happy-nook TM (2015) First Nations perspectives on sea otter conservation in British Columbia and Alaska: insights into coupled human–ocean systems. In: Larson SE, Bodkin JL, VanBlaricom GR (eds) *Sea otter conservation*. Academic Press, Boston, MA, p 301–331
- Sandweiss DH, Andrus CFT, Kelley AR, Maasch KR, Reitz EJ, Roscoe PB (2020) Archaeological climate proxies and the complexities of reconstructing Holocene El Niño in coastal Peru. *Proc Natl Acad Sci USA* 117:8271–8279
- Schaan DP (2016) *Sacred geographies of ancient Amazonia: historical ecology of social complexity*, Vol 3. Routledge, London
- Schroeder D, Chatfield K, Singh M, Chennells R, Herissone-Kelly P (2019) *Equitable research partnerships: a global code of conduct to counter ethics dumping*. Springer, Heidelberg
- Searcy RT, Boehm AB, Weinstock C, Preston CM and others (2022) High-frequency and long-term observations of eDNA from imperiled salmonids in a coastal stream: temporal dynamics, relationships with environmental factors, and comparisons with conventional observations. *Environ DNA* 4:776–789
- Shackeroff JM, Campbell LM, Crowder LB (2011) Social-ecological guilds: putting people into marine historical ecology. *Ecol Soc* 16:52
- Shen Y, Heacock L, Elias J, Hentel KD, Reig B, Shih G, Moy L (2023) ChatGPT and other large language models are double-edged swords. *Radiology* 307:e230163
- Sherwood OA, Lehmann MF, Schubert CJ, Scott DB, McCarthy MD (2011) Nutrient regime shift in the western North Atlantic indicated by compound-specific $\delta^{15}\text{N}$ of deep-sea gorgonian corals. *Proc Natl Acad Sci USA* 108:1011–1015
- Shiraiwa S, Zabalueva O (2021) Museological myths of decolonization and neutrality. In: Bergeron Y, Rivet M (eds) *Proc ICOFOM Symposium: The Decolonisation of Museology: Museums, Mixing, and Myths of Origin*. ICOFOM, Paris, p 203–207
- Sibert EC, Cramer KL, Hastings PA, Norris RD (2017) Methods for isolation and quantification of microfossil fish teeth and elasmobranch dermal denticles (ichthyoliths) from marine sediments. *Palaeontol Electronica* 20: 1–14
- Sidik SM (2022) Weaving Indigenous knowledge into the scientific method. *Nature* 601:285–287
- Silverman H, Ruggles DF (eds) (2009) *Intangible heritage embodied*. Springer, Heidelberg
- Simpson LB (2017) *As we have always done: Indigenous freedom through radical resistance*. U of Minnesota Press, Minneapolis, MN
- Sinclair DJ, Kinsley LP, McCulloch MT (1998) High resolution analysis of trace elements in corals by laser ablation ICP-MS. *Geochim Cosmochim Acta* 62:1889–1901
- Singh GG, Harden-Davies H, Allison EH, Cisneros-Montemayor AM, Swartz W, Crosman KM, Ota Y (2021) Will understanding the ocean lead to 'the ocean we want'? *Proc Natl Acad Sci USA* 118:e2100205118
- Slot M, Winter K (2018) High tolerance of tropical sapling growth and gas exchange to moderate warming. *Funct Ecol* 32:599–611
- Smith NF, Lepofsky D, Toniello G, Holmes K, Wilson L, Neudorf CM, Roberts C (2019) 3500 years of shellfish mariculture on the Northwest Coast of North America. *PLOS ONE* 14:e0211194
- Soga M, Gaston KJ (2018) Shifting baseline syndrome: causes, consequences, and implications. *Front Ecol Environ* 16:222–230
- Spalding AK, Grorud-Colvert K, Allison EH, Amon DJ and others (2023) Engaging the tropical majority to make ocean governance and science more equitable and effective. *npj Ocean Sustainability* 2:8
- Speare KE, Adam TC, Winslow EM, Lenihan HS, Burkepille DE (2022) Size-dependent mortality of corals during marine heatwave erodes recovery capacity of a coral reef. *Glob Change Biol* 28:1342–1358

- Stahl PW (2018) Historical ecology and archaeology in theory and practice. *Antiquity* 92:1677–1679
- Staples TL, Kiessling W, Pandolfi JM (2022) Emergence patterns of locally novel plant communities driven by past climate change and modern anthropogenic impacts. *Ecol Lett* 25:1497–1509
- Starkey DJ, Holm P, Barnard M (eds) (2012) *Oceans past: management insights from the history of marine animal populations*. Earthscan, London
- Stephenson PJ, Grace MK, Akçakaya HR, Rodrigues ASL and others (2019) Defining the indigenous ranges of species to account for geographic and taxonomic variation in the history of human impacts: reply to Sanderson 2019. *Conserv Biol* 33:1211–1213
- Stropp J, Ladle RJM, Malhado ACM, Hortal J and others (2016) Mapping ignorance: 300 years of collecting flowering plants in Africa. *Glob Ecol Biogeogr* 25:1085–1096
- Sturges F (2019) *Humane Society of the United States v. Zinke* (DC CIR. 2017): shifting baselines in the Endangered Species Act. *Harvard Environ Law Rev* 43:225–245
- Sullivan BL, Wood CL, Iliff MJ, Bonney RE, Fink D, Kelling S (2009) eBird: a citizen-based bird observation network in the biological sciences. *Biol Conserv* 142:2282–2292
- Sutton S (2020) The evolving responsibility of museum work in the time of climate change. *Mus Manag Curator* 35: 618–635
- Swanson KL, Sugihara G, Tsonis AA (2009) Long-term natural variability and 20th century climate change. *Proc Natl Acad Sci USA* 106:16120–16123
- Swetnam TW, Allen CD, Betancourt JL (1999) Applied historical ecology: using the past to manage for the future. *Ecol Appl* 9:1189–1206
- Szabó P, Hédl R (2011) Advancing the integration of history and ecology for conservation. *Conserv Biol* 25:680–687
- Szabó P, Kuneš P, Svobodová-Svitavská H, Švarcová MG and others (2017) Using historical ecology to reassess the conservation status of coniferous forests in Central Europe. *Conserv Biol* 31:150–160
- Tanaka KR, Van Houtan KS (2022) The recent normalization of historical marine heat extremes. *PLOS Clim* 1: e0000007
- Taylor C (1992) *Sources of the self: the making of the modern identity*. Harvard University Press, Cambridge, MA
- Thurstan RH (2022) The potential of historical ecology to aid understanding of human–ocean interactions throughout the Anthropocene. *J Fish Biol* 101:351–364
- Toniello G, Lepofsky D, Lertzman-Lepofsky G, Salomon AK, Rowell K (2019) 11,500 y of human–clam relationships provide long-term context for intertidal management in the Salish Sea, British Columbia. *Proc Natl Acad Sci USA* 116:22106–22114
- Trant AJ, Nijland W, Hoffman KM, Mathews DL, McLaren D, Nelson TA, Starzomski BM (2016) Intertidal resource use over millennia enhances forest productivity. *Nat Commun* 7:12491
- Trisos CH, Auerbach J, Katti M (2021) Decoloniality and anti-oppressive practices for a more ethical ecology. *Nat Ecol Evol* 5:1205–1212
- Tuia D, Kellenberger B, Beery S, Costelloe BR and others (2022) Perspectives in machine learning for wildlife conservation. *Nat Commun* 13:792
- Turton D (2011) Wilderness, wasteland or home? Three ways of imagining the Lower Omo Valley. *J East Afr Stud* 5: 158–176
- Van Houtan KS, Kittinger JN (2014) Historical commercial exploitation and the current status of Hawaiian green turtles. *Biol Conserv* 170:20–27
- Van Houtan KS, McClenachan L, Kittinger JN (2013) Seafood menus reflect long-term ocean changes. *Front Ecol Environ* 11:289–290
- Van Houtan KS, Gagne T, Jenkins CN, Joppa L (2020) Sentiment analysis of conservation studies captures successes of species reintroductions. *Patterns* 1:100005
- Wang Y, Pineda-Munoz S, McGuire JL (2023) Plants maintain climate fidelity in the face of dynamic climate change. *Proc Natl Acad Sci USA* 120:e2201946119
- Wells JC, Silva AP, Araújo L, Azevêdo G and others (2019) Empowering communities to identify, treat, and protect their heritage: a cultural landscape case study of the Horto d'El Rey, Olinda, Brazil. In: Fouseki K, Gutormsen T, Swensen G (eds) *Heritage and sustainable urban transformations: deep cities*. Routledge, Abingdon, p 185–207
- West P, Igoe J, King D (2006) Parks and peoples: the social impact of protected areas. *Annu Rev Anthropol* 35: 251–277
- West CF, Etnier MA, Barbeaux S, Partlow MA, Orlov AM (2022) Size distribution of Pacific cod (*Gadus macrocephalus*) in the North Pacific Ocean over 6 millennia. *Quat Res* 108:43–63
- Whitman E, Parisien MA, Thompson DK, Flannigan MD (2019) Short-interval wildfire and drought overwhelm boreal forest resilience. *Sci Rep* 9:18796
- Wickham SB, Augustine S, Forney A, Mathews DL, Shackelford N, Walkus J, Trant AJ (2022) Incorporating place-based values into ecological restoration. *Ecol Soc* 27:32
- Wilkinson MD, Dumontier M, Aalbersberg IJ, Appleton G and others (2016) The FAIR Guiding Principles for scientific data management and stewardship. *Sci Data* 3: 160018
- Witter R, Satterfield T (2019) The ebb and flow of Indigenous rights recognitions in conservation policy. *Dev Change* 50:1083–1108
- Witze A (2024) Geologists reject the Anthropocene as Earth's new epoch — after 15 years of debate. *Nature* 627: 249–250
- Wolfe AP, Hobbs WO, Birks HH, Briner JP and others (2013) Stratigraphic expressions of the Holocene–Anthropocene transition revealed in sediments from remote lakes. *Earth Sci Rev* 116:17–34
- Wolverton SJ, Nagaoka L, Rick TC (2016) *Applied zooarchaeology: five case studies*. Eliot Werner Publications, Clinton Corners, NY
- Xu L, Liu X, Wu L, Sun L, Zhao J, Chen L (2016) Decline of recent seabirds inferred from a composite 1000-year record of population dynamics. *Sci Rep* 6:35191
- Yasuoka H (2013) Dense wild yam patches established by hunter-gatherer camps: beyond the wild yam question, toward the historical ecology of rainforests. *Hum Ecol* 41: 465–475
- Zhang DD, Brecke P, Lee HF, He YQ, Zhang J (2007) Global climate change, war, and population decline in recent human history. *Proc Natl Acad Sci USA* 104: 19214–19219
- Zimmerman SR, Wahl DB (2020) Holocene paleoclimate change in the western US: the importance of chronology in discerning patterns and drivers. *Quat Sci Rev* 246: 106487

Appendix. List of author affiliations

Loren McClenachan^{1,2}, Torben Rick³, Ruth H. Thurstan⁴, Andrew Trant⁵, Peter S. Alagona⁶, Heidi K. Alleway⁷, Chelsey Armstrong⁸, Rebecca Bliege Bird⁹, Nadia T. Rubio-Cisneros^{10,11}, Miguel Clavero¹², André C. Colonese¹³, Katie Cramer¹⁴, Ancilleno O. Davis¹⁵, Joshua Drew¹⁶, Michelle M. Early-Capistrán¹⁷, Graciela Gil-Romera¹⁸, Molly Grace¹⁹, Marco B. A. Hatch²⁰, Eric Higgs², Kira Hoffman^{21,22}, Jeremy B. C. Jackson²³, Antonieta Jerardino²⁴, Michelle J. LeFebvre²⁵, Heike K. Lotze²⁶, Ryan S. Mohammed²⁷, Naia Morueta-Holme²⁸, Catalina Munteanu²⁹, Alexis M. Mychajliw³⁰, Bonnie Newsom³¹, Aaron O'Dea^{32,33}, Daniel Pauly³⁴, Péter Szabó^{35,36}, Jimena Torres^{37,38}, John Waldman³⁹, Catherine West⁴⁰, Liqiang Xu⁴¹, Hirokazu Yasuoka⁴², Philine S. E. zu Ermgassen⁴³, Kyle S. Van Houtan⁴⁴

¹Department of History, University of Victoria, Victoria, BC V8W 2Y2, Canada

²School of Environmental Studies, University of Victoria, Victoria, BC V8W 2Y2, Canada

³Department of Anthropology, National Museum of Natural History, Smithsonian Institution, Washington, DC 20560, USA

⁴Centre for Ecology and Conservation, University of Exeter, Penryn TR10 9FE, UK

⁵School of Environment, Resources and Sustainability, University of Waterloo, Waterloo, ON N2L 3G1, Canada

⁶Environmental Studies Program, University of California, Santa Barbara, CA 93106, USA

⁷The Nature Conservancy, Arlington, VA 22203, USA

⁸Indigenous Studies, Simon Fraser University, Burnaby, BC V5A 1S6, Canada

⁹Department of Anthropology, Pennsylvania State University, State College, PA 16801, USA

¹⁰Facultad de Ciencias Biológicas, Universidad Autónoma de Nuevo León, San Nicolás de los Garza 66450, Mexico

¹¹Mar Sustentable Ciencia y Conservación, Monterrey 66450, Mexico

¹²Estación Biológica de Doñana, Consejo Superior de Investigaciones Científicas, Sevilla 41092, Spain

¹³Department of Prehistory and Institute of Environmental Science and Technology, Universitat Autònoma de Barcelona, Barcelona 08193, Spain

¹⁴Center for Biodiversity Outcomes, Arizona State University, Tempe, AZ 85287, USA

¹⁵The Bahamas National Trust, Nassau, The Bahamas

¹⁶Department of Environmental Biology, State University of New York College of Environmental Science and Forestry, Syracuse, NY 13210, USA

¹⁷Hopkins Marine Station, Stanford University, Pacific Grove, CA 93950, USA

¹⁸Instituto Pirenaico de Ecología, Consejo Superior de Investigaciones Científicas, Zaragoza 50059, Spain

¹⁹Department of Biology, University of Oxford, Oxford OX1 3RB, UK

²⁰Environmental Sciences, Western Washington University, Bellingham, WA 98225, USA

²¹Faculty of Forestry, University of British Columbia, Vancouver, BC V6T 1Z4, Canada

²²Bulkley Valley Research Centre, Smithers, BC V0J 2N0, Canada

²³Division of Paleontology, American Museum of Natural History, New York, NY 10024, USA

²⁴Department of Anthropology & Archaeology, University of South Africa, Pretoria 0028, South Africa

²⁵Florida Museum of Natural History, University of Florida, Gainesville, FL 32611, USA

²⁶Department of Biology, Dalhousie University, Halifax, NS B3H 4R2, Canada

²⁷Department of Biological Sciences, Auburn University, Auburn, AL 36849, USA

²⁸Center for Macroecology, Evolution and Climate, Globe Institute, University of Copenhagen, 2100 Copenhagen, Denmark

²⁹Faculty of Environment and Natural Resources, University of Freiburg, 79106 Freiburg, Germany

³⁰Department of Biology and Program in Environmental Studies, Middlebury College, Middlebury, VT 05753, USA

³¹Department of Anthropology, University of Maine, Orono, ME 04469, USA

³²Smithsonian Tropical Research Institute, Balboa 0843-03092, Republic of Panamá

³³Sistema Nacional de Investigación, Secretaría Nacional de Ciencia Tecnología e Innovación, Panamá City 0801, Republic of Panamá

³⁴Sea Around Us, Institute for the Oceans and Fisheries, University of British Columbia, Vancouver, BC V6T 1Z4, Canada

³⁵Department of Vegetation Ecology, Institute of Botany, Czech Academy of Sciences, 252 43 Průhonice, Czech Republic

³⁶Department of Environmental Studies, Faculty of Social Sciences, Masaryk University, 252 43 Průhonice, Czech Republic

³⁷Cape Horn International Center, Universidad de Magallanes, Cabo de Hornos, Chile

³⁸Millennium Nucleus Program, UPWELL Foundation, Chile

³⁹Queens College and Graduate School, City University of New York, Flushing, NY 11367, USA

⁴⁰Department of Anthropology and Archaeology Program, Boston University, Boston, MA 02215, USA

⁴¹School of Resources and Environmental Engineering, Hefei University of Technology, Anhui 230036, PR China

⁴²Center for African Area Studies, Kyoto University, Sakyo, Kyoto 606-8304, Japan

⁴³Changing Oceans Group, School of Geosciences, University of Edinburgh, Edinburgh EH8 9XP, UK

⁴⁴Nicholas School of the Environment, Duke University, Durham, NC 27710, USA