

Developing an eDNA toolkit to quantify broadcast spawning events of the sea scallop *Placopecten magellanicus*: moving beyond fertilization assays

Skylar R. Bayer^{1,3,*}, Peter D. Countway², Richard A. Wahle¹

¹University of Maine, School of Marine Sciences, Darling Marine Center, 193 Clark's Cove Road, Walpole, ME 04573

²Bigelow Laboratory for Ocean Sciences, 60 Bigelow Drive, East Boothbay, ME 04544

³Present Address: NOAA NEFSC Milford Laboratory, 212 Rogers Avenue, Milford, CT 06460

ABSTRACT: Broadcast spawners release their gametes into the water column for 'chance' fertilization events. However, detection of such events in near real-time is extremely difficult, but needed to improve fisheries and conservation management practices. It is practically impossible to distinguish the gametes of many species by microscopy; therefore, DNA-based techniques are preferable to detect and quantify gametes from field-collected plankton samples. We developed a quantitative PCR (qPCR) approach to detect and quantify broadcast spawning events in marine environmental DNA (eDNA) samples. We applied this approach to a commercially valuable broadcast spawning bivalve species, the sea scallop *Placopecten magellanicus*. Our approach includes (1) sequencing the internal transcribed spacer (ITS) region, (2) developing a novel species-specific probe and primer set, (3) testing the probe and primer set on a dilution series of sea scallop sperm to quantify the relationship between gamete abundance and DNA copy number, and (4) conducting dockside field tests of our method on plankton samples adjacent to naturally spawning sea scallops. Quantitative PCR revealed a clear relationship between DNA copy number and *P. magellanicus* sperm cell abundance, indicating that this method is reliable for detecting sperm release by male scallops during spawning events. Plankton samples collected during the scallop spawning season revealed spikes of scallop eDNA in both the <20 µm (sperm) and >20 µm (possible eggs) particle size-fractions. This method holds great potential to provide more efficient estimates of the timing, magnitude, and spatial scale of reproductive events than conventional methods for a wide range of broadcast spawners.

KEY WORDS: Sea scallop · *Placopecten magellanicus* · Reproductive ecology · eDNA · ITS region · Gamete detection · Broadcast spawning

Resale or republication not permitted without written consent of the publisher

1. INTRODUCTION

Marine organisms spanning multiple phyla broadcast gametes into the sea where successful fertilization depends on having high concentrations of viable sperm. High-density spawning aggregations of adults can ensure high levels of fertilization success (the percentage of eggs fertilized) before game-

tes become diluted (Allee 1931, Levitan et al. 1992, Levitan & Petersen 1995, Levitan & Young 1995, Gaudette et al. 2006). Empirical data on fertilization success in natural broadcast spawner populations are relatively rare and generally qualitative (Levitan & Sewell 1998). To date, field studies on fertilization dynamics have been conducted on a relatively small set of experimentally tractable free spawning inver-

*Corresponding author: skylar.bayer@noaa.gov

tebrates and fishes: including sea urchins *Strongylocentrotus droebachiensis* (Wahle & Peckham 1999), bluehead wrasse *Thalassoma bifasciatum* (Petersen et al. 1992, Warner et al. 1995), hard corals (*Montipora digitata*, *Favites pentagona*, *Platygyra sinensis*, Oliver & Babcock 1992; *Acanthaster planci*, Babcock & Mundy 1992), soft corals *Briareum asbestinum* (Brazeau & Lasker 1992), and sea cucumbers (*Bohadschia argus*, *Eupta godeffroyi*, *Stichopus variegatus*, *Holothuria coluber*, *Actinopyga lecanora*, *Bohadschia graffei*; Babcock et al. 1992). Together they suggest that fertilization success can be quite variable and frequently well below saturating levels, especially at low adult densities and conditions promoting rapid gamete dilution.

These results underscore a long-standing concern that commercially exploited fishes and invertebrates with this mode of reproduction may be especially vulnerable to recruitment failure when populations are depleted (Myers et al. 1995, Petersen & Levitan 2001, Rowe et al. 2004). Aggregating behavior has been observed specifically during the spawning season. Sedentary or slow-moving broadcast spawners that form spawning aggregations, such as the abalone *Haliotis kamtschatkana* (Seamone & Boulding 2011) and scallop (*Chlamys asperrima* and *Chlamys bifrons*, Styan & Butler 2003; *Pecten fumatus*, Mendo et al. 2014), may be particularly vulnerable to the effects of fishing activity that target these aggregations.

Field experiments with experimentally tractable species that are relatively sedentary and easy to induce to spawn have contributed considerably to understanding the importance of sperm limitation in the wild, but they also illustrate the challenges of measuring fertilization *in situ*. Some investigators have collected naturally spawned gametes in the wild using various pumps, filters, suction samplers and large bags (e.g. Levitan 1991, Petersen et al. 1992, 2001, Warner et al. 1995, Styan 1997). These methods may bias estimates of fertilization by artificially exposing eggs to high concentrations of sperm for longer than they would normally experience (Levitan 1995). Models of fertilization kinetics predict that most sperm and egg collisions occur within the first few seconds of exposure, and that experimental artifacts biasing the measure of fertilization success may come from artificially altering the duration of exposure (Vogel et al. 1982, Styan 1998b). Several field studies have deployed freshly spawned, unfertilized eggs in synthetic mesh (e.g. nylon) chambers that retain the eggs but are permeable to sperm (Levitan et al. 1992, Levitan & Young 1995, Wahle & Peckham 1999, Gaudette et al. 2006, Bayer et al.

2016). Such chambers inevitably alter natural transport of gametes, and can produce experimental artifacts by impeding flow while keeping eggs stationary, which increases the amount of water—and therefore the amount of sperm—encountered by eggs (Levitan 1991, Levitan et al. 1992). Nonetheless, these methods have been recommended for eggs that are barely visible and are found at low concentrations (Styan 1997).

Molecular techniques to identify and quantify species-specific DNA from sea water samples through genetic markers have been applied extensively to eukaryotic microbes (Countway & Caron 2006, Moorthi et al. 2006, Garneau et al. 2011), marine viruses (Hewson et al. 2011, Matteson et al. 2013) and diverse groups of bacteria and archaea (Suzuki et al. 2000, Schwabach & Fuhrman 2005, Ahlgren & Rocop 2012). These same molecular techniques have been used to identify and quantify metazoan DNA including that of invertebrate larvae (abalone, Vadopalas et al. 2006; sea lice, McBeath et al. 2006; decapods, Pan et al. 2008), marine fin fish embryos (Taylor et al. 2002, Fox et al. 2005) and freshwater fin fish spawning events (perch, Bylemans et al. 2017). Information gained from these studies have helped estimate larval dispersal (McBeath et al. 2006) and inform finfish stock assessments (Fox et al. 2005).

The sea scallop *Placopecten magellanicus* is a sedentary, gonochoristic, broadcast spawner found in the coastal and shelf waters of the Northwest Atlantic. Rescued from the brink of widespread depletion through a strategy of controls on fishing effort and closures of fishing grounds, the sea scallop is considered one of few existing success stories in fishery management (Hart & Rago 2006). While key components of recruitment, such as adult gamete production (Langton et al. 1987), larval transport (Tremblay et al. 1994, Tian et al. 2009), juvenile mortality (Wong & Barbeau 2003) and patchiness (Carey et al. 2013), have been subjects of intensive study, the recent literature reflects a growing recognition of how little empirical information is available on the role of spawning and fertilization dynamics as a potentially critical step in the recruitment process for scallops (Stokesbury 1999, 2012, Smith & Rago 2004). While information on the timing of spawning events can be gathered from gonad indices (Langton et al. 1987, Bonardelli et al. 1996, Thompson et al. 2014, Bayer et al. 2016), these data are limited in temporal resolution to days and weeks and can only be generalized over broad areas, thereby compromising the detection and measurement of finer-scale local spawning events. While field-deployed time-integrated fertil-

ization assays have begun to open a window on fine-scale spawning dynamics in sea scallops (Bayer et al. 2016, 2018), they are time- and labor-intensive and are prone to experimental artifacts, limiting their broad practical application.

Molecular techniques used to detect invertebrate larvae and eukaryotic plankton are well suited for the detection of sea scallop sperm presence and abundance from sea water samples. The internal transcribed spacer (ITS) region, located between the small (18S) and large (28S) subunits of the rRNA gene, is a useful locus for eDNA-based studies of marine eukaryotes. For example, previous studies have targeted the ITS region of harmful microalgae (Andree et al. 2011, Vandersea et al. 2017), fungi (Op De Beeck et al. 2014), parasites (Reece et al. 2008) and bivalves (Insua et al. 2003, Vierna et al. 2010, Lazoski et al. 2011, Salvi et al. 2014). Wang et al. (2006) found that the ITS region was a reliable genetic marker for species identification among 12 commercially harvested bivalves. The ITS region is evolutionary conserved within a species, but is diverse among species, even those closely related within a group such as Pectinidae (Insua et al. 2003). This study describes development and application of a real-time quantitative PCR assay for high-throughput detection and quantification of sea scallop sperm in seawater samples. The steps we took in developing this method included (1) sequencing a unique and reliable region, the ITS, of the sea scallop genome, (2) developing a qPCR probe and primer set to detect a diagnostic DNA fragment within this region, (3) testing the developed probe and primer set on a dilution series of scallop sperm and eggs to quantify the relationship between gamete abundance and DNA copy number, and (4) dockside field tests to evaluate the method with plankton samples collected adjacent to naturally spawning scallops. Our novel approach revealed dynamic trends in scallop eDNA abundance that coincided with the timing of an expected spawning event, and also captured finer-scale details in terms of the timing and magnitude of spawning than would have been possible using traditional approaches.

2. MATERIALS AND METHODS

2.1. DNA extraction, cultures, cloning and sequencing

We sampled *Placopecten magellanicus* mantle tissue of individuals collected from the Damariscotta River estuary using a minimally invasive approach,

and stored this material at -20°C before extraction using the E.Z.N.A.[®] Mollusc DNA Kit (Omega Bio-Tek), following the manufacturer's protocol. DNA was quantified using a Qubit fluorometer and the Broad Range (BR) Qubit detection reagents. PCR for molecular cloning of *P. magellanicus* DNA was performed using ~ 10 ng of purified scallop DNA per reaction and 500 nM each of the eukaryotic primers Euk-B Forward: 5'-GTA GGT GAA CCT GCA GAA GGA TC-3' (Medlin et al. 1988) and NLR204: 5'-ATA TGC TTA ART TCA GCG GGT-3' (Van der Auwera et al. 1994) to amplify the entire ITS region between the small-subunit ribosomal RNA (rRNA) and large-subunit rRNA genes, including ITS1, 5.8S, and ITS2. A total of 6 PCRs were set up, using 25 μl of GoTaq[®] G2 Hot Start Green Master Mix (Promega), primers and DNA as noted above, and molecular grade water to achieve a total volume of 50 μl for each reaction. PCR was performed on a Bio-Rad C1000 PCR machine using a touchdown thermal protocol consisting of initial denaturation at 95°C for 2 min, followed by a touchdown phase that included 10 cycles of denaturation at 95°C for 30 s, annealing at 65 to 55°C for 30 s (decreasing by 1°C per cycle), and extension at 72°C for 90 s. An additional 25 cycles of denaturation at 95°C for 30 s, annealing at 55°C for 30 s, and extension at 72°C for 90 s completed the amplification. The amplified DNA was run on a 1% agarose gel to isolate an ~ 750 bp PCR product spanning the entire ITS region of *P. magellanicus*. The ~ 750 bp PCR product was excised from the gel and purified using the Wizard[®] SV Gel and PCR Clean-Up kit (Promega). PCR products were ligated into the pCR4-TOPO[®] cloning vector and cloned into One Shot[®] TOP-10 chemically competent cells following the manufacturer's protocols included with the TOPO[®] TA Cloning[®] Kit for Sequencing (ThermoFisher Scientific). ITS clones were plated and grown overnight on LB plates with Kanamycin ($50 \mu\text{g ml}^{-1}$). Individual colonies were picked from the plates and grown overnight in Terrific Broth broth with Kanamycin ($50 \mu\text{g ml}^{-1}$) prior to extraction of plasmid DNA using the Zippy[™] Plasmid Miniprep Kit (Zymo Research). Eight of the ITS clones were selected at random and sent to Eurofins Genomics (Louisville, Kentucky) for Sanger DNA sequencing using T3 and T7 sequencing primers to obtain full-length ITS sequences. *P. magellanicus* ITS sequences were submitted to GenBank under accession numbers: MG744618-MG744625. Two micrograms of plasmid DNA from each of the clones was linearized with the *NotI*-HF[®] restriction enzyme (New England Biolabs) in 12 h reactions, prior to band-isolation on a

0.8% TAE gel and purification with the Wizard® SV gel kit (Promega). Linearized plasmid DNA was subsequently used as a DNA standard for the *P. magellanicus* qPCR assay.

2.2. Probe and primer design

We compared the 8 *P. magellanicus* ITS sequences (amplicons, Table 1) to all sequences in the GenBank nr database via BLAST analysis and confirmed that they were homologous to ITS sequences from other scallop species. To the best of our knowledge, these sequences are the first for *P. magellanicus* that span the entire ITS region. Using our newly sequenced ITS clones, we developed a Taqman® minor groove binding (MGB) qPCR assay (Holland et al. 1991, Heid et al. 1996) to quantify *P. magellanicus* DNA in environmental samples. The 8 clone sequences were aligned in Geneious (v.9.1) using MUSCLE (Edgar 2004) and then imported into AlleleID (v.7.83) to design a novel DNA probe, forward, and reverse qPCR primers. The resulting primers and probe were compared to all sequences in GenBank using BLAST (Altschul et al. 1997), to minimize the chance of false positives due to matches with DNA from non-target organisms in seawater samples. The analysis revealed that the primers and probe for *P. magellanicus* ITS were highly specific for the targeted species, revealing no significant matches to other marine species. The forward primer (Pmag_282F: 5'-CAG CCT GAA ATC GAG ATG-3') was located at positions 282 through 299 and the reverse primer (Pmag_492R: 5'-CAT GCA TAC AGT TGC GAT A-3') at positions 474 through 492 relative to our *P. magellanicus* ITS sequences, resulting in a qPCR amplicon of 211 bp in length (Table S1 in the Supplement at www.int-res.com/articles/suppl/m621p127_supp.pdf). The probe (Pmag_304F:5'-[6FAM] TAT TAT CAC TCT AAG

CGG[NFQ]-3') was located very close to the forward primer at positions 304 through 321, and was labelled on the 5'-end with the fluorescent reporter dye 6-carboxyfluorescein (FAM) and on the 3'-end with a proprietary non-fluorescent quenching (NFQ) molecule (ThermoFisher Scientific) to suppress FAM fluorescence in the absence of amplification (Table 1). The probe was modified with a DNA MGB to increase its melting temperature. The FAM molecule is released from the probe during DNA amplification by the 5'-exonuclease activity of Taq DNA polymerase, causing emission of a fluorescence signal that is captured by a detector at the end of the combined annealing/extension step. The cumulative increase in fluorescence is proportional to the amount of amplified DNA, and the probe-based TaqMan assay ensures that the fluorescence is due to amplification of the specific DNA target only (Holland et al. 1991).

2.3. Calibration and testing of qPCR method

Copy numbers per microliter of the cloned PCR products were calculated for each of the 8 *P. magellanicus* ITS clones based on their specific DNA concentrations and the exact nucleotide composition of each PCR product plus cloning vector. Each of the ITS clones had a slightly different DNA sequence, presumably due to the presence of multiple copies of the ITS region in the sea scallop genome and potentially due to PCR-induced errors during DNA replication. However, many of the nucleotide polymorphisms across the full-length ITS region did not appear to be random and displayed the same polymorphisms in >1 sequence, suggesting that the variation was not artifactual. Within the 211 bp region targeted by our qPCR assay, there were only 4 polymorphic nucleotides among the 8 ITS clones located at 3 polymorphic sites. Two of the sequences displayed single polymorphisms at 2 sites, while 2 sequences had the same polymorphism at a third site. None of the polymorphisms occurred within the regions where the probe or primers would bind, thus minimizing the effect of sequence variation on the qPCR assay. To account for some of the potential variability of the qPCR assay due to slight variations in sequence composition of DNA targets in natural samples, equal copy numbers of each of the 8 clones were mixed to produce a composite linear

Table 1. TaqMan probe and primer sequences: properties including sequence, length, melting temperature (T_m) and GC base content of *Placopecten magellanicus*-specific TaqMan probe and primers

Name	Sequence (5'→3')	Length (nucleotides)	T_m (°C)	% GC
Pmag_304F (MGB-probe)	TAT TAT CAC TCT AAG CGG	18	68	38.9
Pmag_282F (primer)	CAG CCT GAA ATC GAG ATG	18	59.1	50
Pmag_492R (primer)	CAT GCA TAC AGT TGC GAT A	19	59.2	42.1

DNA standard with a final concentration of 3.7×10^9 copies μl^{-1} . The DNA standard was serially diluted to create a standard curve spanning 7 orders of magnitude (i.e. 10^{-2} to 10^{-9}). Quantitative estimates of scallop DNA in environmental samples must be considered minimal estimates due to the potential for inhibition of the qPCR assay by natural (and unknown) inhibitor molecules and/or competition for probe and primers by non-target (and unknown) DNA templates present in our eDNA extracts. Inhibition of the qPCR assay by unknown molecules in the DNA extract should have been minimal due to our use of the proprietary Inhibitor Removal Technology within the Qiagen PowerWater extraction kit. Given the small size of our qPCR target (211 bp), ~96% of the DNA in each of the dilution-series standards was non-target DNA from the pCR4-TOPO cloning vector and could be considered 'background DNA' to approximate any 'sample matrix' effects.

All reagents, samples, and standards were prepared on ice prior to thermal cycling. Reactions of DNA standards were prepared as follows: 3 μl aliquot of each diluted ITS standard, 10 μl of 2' Prime-Time® Gene Expression Master Mix (Integrated DNA Technologies), 1 μl each of the forward and the reverse primers (500 nM final concentration of each), 0.5 μl of the probe (250 nM final concentration), and 4.5 μl of deionized water. All samples, standards and no-template controls were loaded in triplicate on a white-well PCR plate (Bio-Rad) and amplified on a Bio-Rad CFX96 real-time PCR system. Thermal cycling consisted of an initial denaturation step of 3 min at 95°C, followed by 40 cycles of 95°C for 15 s and 61°C for 30 s. Real-time relative fluorescence unit (RFU) values were collected after the combined annealing/extension step.

2.4. Cross-reactivity testing

Although our *P. magellanicus* qPCR assay was designed to be species-specific and was tested against closely related taxa *in silico*, we also conducted *in vivo* cross-reactivity testing against 2 bivalve species common to the Gulf of Maine. To test for cross-reactivity of the probe and primers, we collected DNA from the eastern oyster *Crassostrea virginica* and the blue mussel *Mytilus edulis*. We obtained an archived DNA sample from *C. virginica* that had been collected from Jones Cove (Damariscotta River, ME) and extracted by Marquis et al. (2015). Blue mussels were collected from the dock at the University of Maine's Darling Marine Center

(Damariscotta River, ME, USA), and DNA was extracted from the mantle tissue of 1 individual following the same protocol used for *P. magellanicus* described above. PCR, cloning, and sequencing of the ITS region for *M. edulis* and *C. virginica* also followed the procedures described above. Triplicate qPCR reactions for each of the non-target bivalve species were conducted using 10^{-3} and 10^{-6} dilutions of each of the cloned, non-target DNA templates. Three ITS clones from each non-target species were sequenced to confirm their identity and ensure non-reactivity with the *P. magellanicus* qPCR primers and probe (Table S2, GenBank accession numbers MG744626-MG744631).

2.5. Collection of scallop gametes

Male sea scallops, that were originally collected from the Damariscotta River estuary and kept in hanging nets on docks at the Darling Marine Center, were selected for spawning during the first week of September 2016. Spawning tables (92 × 122 cm) were filled 15 cm deep with filtered (1 μm) UV-sterilized sea water and cooled to 10°C. Male scallops were transferred to spawning tables from ambient (~16°C) unfiltered seawater and induced to spawn via a cold shock as in Bayer et al. (2016). From these, 250 ml of fresh, concentrated sperm cells were collected in a 250 ml, acid-washed container and serially diluted twice before filtering onto 0.2 μm , 47 mm diameter Supor filters (Whatman) to simulate collection of a plankton-based eDNA sample. Filters were rolled gently, placed in cryo-vials and stored at -20°C until DNA extraction. This dilution series preparation and filtration was repeated 5 times, with each initial sample collected from ambient sperm concentrations within the spawning table, and subsamples collected into 1.5 ml vials, and preserved in 4% formalin for sperm cell counts by microscopy. Three subsamples of sperm cells were counted with a haemocytometer using standard counting protocols (e.g. Wahle & Peckham 1999, Andersen & Thronsdon 2004, Gaudette et al. 2006).

2.6. Sperm dilution series

DNA was extracted from filtered-collected and frozen sperm cell samples using the PowerWater DNA extraction and purification kit (Qiagen) with a slight modification of the sample lysis step. Briefly, frozen Supor filters were transferred to 5 ml Power-

Water bead tubes containing pre-loaded garnet particles. One ml of heated (55°C) PW1 lysis buffer was added to the bead tubes and heated at 55°C for 3 to 5 min. The 5 ml tubes were transferred to a 5 ml sample-tube adapter (MoBio) and loaded onto a MM400 Retsch Mixer Mill for sample bead-beating. Two rounds of bead-beating were conducted for 5 min each, at an oscillation frequency of 20 s⁻¹, with 180° rotation of the tube adapter between the 5 min bead-beating sessions. Additional sample processing followed the Qiagen PowerWater protocol exactly. TaqMan qPCR was performed on the extracted samples with the *P. magellanicus* probe and primers described above using 3 µl aliquots of each purified DNA sample in triplicate, with 3 replicates of a filtered, UV-sterilized seawater control. Using the standard curve that was generated from the linearized-plasmid dilution series, we translated the Cq values (qPCR threshold cycle number) resulting from cell-based DNA samples to gene copy numbers to investigate the relationship between gene copy number and cell abundance.

2.7. Dockside experiment

All scallops used for the dockside experiment were collected by SCUBA dive surveys in July 2017 from the Damariscotta River estuary near Bigelow Laboratory for Ocean Sciences. Specimens were kept in tanks at the University of Maine's Darling Marine Center until the start of the experiment at Bigelow Laboratory. To independently monitor spawning activity over the duration of our dockside experiments, we tracked gonadosomatic indices (GSI) in a small (<50 individuals) population (i.e. Bayer et al. 2016, 2018) maintained in flowing seawater tanks at the Darling Marine Center, located ~8 km upriver from the Bigelow dock. GSIs were monitored as a general indicator of the onset and end of the spawning season. We sampled between 6 and 10 individuals (with equal sex ratios) 4 times during the spawning season with two 3 wk intervals between sampling events.

Two lantern nets were suspended ~1 m apart under the dock at Bigelow Laboratory for Ocean Sciences, on the lower portion of the Damariscotta River estuary. One net contained 2 females and 9 males, and the other contained 5 males and 5 females, for a total population of 21 individuals. The scallops were spread out over 4 levels of net as in Bayer et al. (2016).

We collected seawater samples with a suction hose pumping at a rate of 15 l min⁻¹ adjacent to the nets.

The end of the sampling hose was deployed <10 cm from the edge of a lantern net, approximately halfway down (~0.5 m) the length of the net. We let the plankton pump run for at least 60 s to rinse the sample hose, then rinsed each of three 6 l carboys before filling them completely. We sampled daily during different phases of the lunar cycle: 1 to 2 times per week during non-target weeks, and more intensively during the week between the full and third quarter moons, based on our previous observations of spawning during that time period (Bayer pers. obs.). On each sampling day, we collected water within 2 h of low or high slack tide to maximize uptake of potential cells downstream from the lantern nets. Water samples were size-fractionated to test for scallop eDNA in particles > 20 µm (inclusive of scallop eggs) and those ranging from <20 to 0.2 µm (inclusive of scallop sperm cells). The 6 l samples were gravity-filtered through 20 µm Nylon mesh filters installed in 47 mm, in-line filter holders (Pall Laboratory) for extraction of DNA from particles > 20 µm. One liter of the <20 µm filtrate was subsequently filtered through a 0.2 µm Supor filter for DNA extraction of particles <20 µm, the size-fraction that included scallop sperm cells. All filters were frozen, and DNA was extracted using the PowerWater DNA kit (Qiagen) as described above.

After running samples through the scallop qPCR protocol, we examined the time-series for noticeable increases (peaks) in gene copy number. For samples < 20 µm, we assumed that all scallop DNA was attributed to sperm and converted gene copy number to sperm cells based on the sperm dilution series described above. In the laboratory, we were unable to develop a linear relationship between cell counts and gene copy numbers for eggs despite sampling freshly spawned scallop eggs that were well-mixed prior to dilutions, filtering, and subsequent DNA extraction. We suspect that the method we developed for sperm cell counts and DNA extraction may not be appropriate for eggs.

2.8. Phylogenetic reconstruction

A total of 14 full-length bivalve species ITS sequences were generated for target (*P. magellanicus*, n = 8) and non-target species (*C. virginica*, n = 3, and *M. edulis*, n = 3). These sequences were combined with other bivalve ITS sequences from GenBank (Benson et al. 2013) to generate a phylogenetic tree within the Geneious v9 software package, using the MUSCLE sequence aligner (Edgar 2004) and the

Bayesian phylogenetic inference program, MrBayes v3.2 (Huelsenbeck & Ronquist 2001). The initial MUSCLE alignment was refined by manually trimming the non-overlapping regions at the 5' and 3' ends, and was subsequently realigned, resulting in an alignment spanning 1635 nucleotide positions (see the Supplement Data file, www.int-res.com/articles/suppl/m621p127_supp/). The aligned sequences were imported into PAUP* v4.0a164 (Swofford 2003) and processed with the MrModelblock file (available from <https://github.com/nylander/MrModeltest2/>) to generate a 'mrmodel.scores' input file for MrModeltest v2.4 (Nylander 2004). The result of MrModeltest indicated that the best nucleotide substitution model for our DNA alignment was the 'general time reversible' (GTR) nucleotide substitution model with rate variation set to 'invgamma' (+I +G), indicating a gamma distribution of nucleotide substitution rates with a proportion of invariable nucleotide sites. An ITS sequence from the marine gastropod *Coralliophila meyendorffii* (Oliveiro & Mariottini 2001) served as the outgroup for this phylogenetic tree.

3. RESULTS

3.1. *Placopecten magellanicus* qPCR calibration and cross-reactivity tests

Real-time qPCR showed a clear logistical relationship between Cq values from cell extracts and gene copy numbers using the probe and primers designed from the *P. magellanicus* ITS region (Fig. 1). The lowest limit of quantification (LOQ) with reproducible amplification was ~11 gene copies per reaction, using 3 μ l of the most dilute DNA standard (10^{-9}) in a 20 μ l reaction volume. The limit of detection, which accounts for all of the factors that might impact the detection of a qPCR target (Forootan et al. 2017), was not determined empirically, given that a single scallop gamete (the basic ecological unit in our study) likely has more copies of the ITS region than the observed LOQ for this assay. The efficiency of the *P. magellanicus* TaqMan qPCR assay was generally in the range of 92 to 96% for all sample runs. No-template control Cq values were

>38, indicating very little background contamination, suggesting that the vast majority of the signal from the cell-based samples was due to the extracted DNA from *P. magellanicus* sperm cells. The UV sterilized seawater sample, serving as a background sample to accompany the sperm dilution series, had a Cq value of ~31 (~100 gene copies). All samples in the sperm dilution series had Cq values ≤ 20 (e.g. ≥ 760400 gene copies), indicating background DNA signals of ~100 gene copies impacted the dilution series samples only minimally. After translating Cq values from the standard curve to gene copy numbers, the sperm dilution series indicated a strong relationship between gene copy number and cell counts by microscopy (Fig. 2). We observed ~1 ITS gene copy per 4 to 5 sperm cells consistently across samples.

We tested our *P. magellanicus* qPCR assay against eastern oyster *Crassostrea virginica* and blue mussel




Fig. 1. Real-time PCR cycle number (Cq) and gene copy number for dilution series of standard clonal *P. magellanicus* ITS regions




Fig. 2. Cell count of *P. magellanicus* sperm and resulting gene copy number from 5 dilution series (red, blue, green, purple and black filled circles)

Mytilus edulis ITS clones that were generated from Damariscotta River specimens. These 2 bivalve species were selected as co-occurring broadcast spawners that could potentially contribute DNA to our samples. Real-time qPCR demonstrated no cross-reactivity of the *P. magellanicus* probe and primers to eastern oyster or blue mussel ITS clones.

3.2. Dockside experiment

In 6 l of pumped seawater, we detected millions of scallop gene copies from samples of cells < 20 μm and thousands of gene copies from cells > 20 μm . Our dockside time-series, consisting of the 1 l, <20 μm samples, indicated possible spawning events on 28 and 29 August and 27 September (Fig. 3a). The scallop eDNA signal in the >20 μm size-fraction was high on 28 and 29 August, and 1 additional sample was very high on 16 September (Fig. 3b). A nearby *P. magellanicus* population at the Darling Marine Center that we monitored for changes in GSI revealed a reduction in male and female gonad size between 19 August and 9 October (Fig. 3c), consistent with spawning activity over the period that we conducted dockside sampling.

3.3. Phylogenetic reconstruction

P. magellanicus ITS sequences varied from 758 to 765 nucleotides in length, and none of the 8 sequences were identical, suggesting a high degree of intragenomic variability and multiple copies of the rRNA operon within the scallop genome. Despite this high variability, the *P. magellanicus* primers and probe were designed to target a conserved region that would amplify all of the known sequence variants that we identified. We found a single, non-annotated *P. magellanicus* cDNA sequence in GenBank (accession: GADG01000430.1) that overlapped partially with our full-length scallop ITS sequences (Pairett & Serb 2013). This cDNA sequence was too short to include in the phylogenetic reconstruction but contained the entire region targeted by our scallop qPCR assay, revealing 100% sequence identity to our newly designed probe and primers (see Fig. S1 in the supplement at www.int-res.com/articles/suppl/m621p127_supp.pdf). All *P. magellanicus* sequences were most similar to ITS sequences from other species of scallops (Fig. 4). Blue mussel and Eastern oyster ITS sequences were most similar to other ITS sequences




Fig. 3. Time series of (a) estimated sperm (cells < 20 μm) abundance and (b) gene copy number from cells > 20 μm from plankton pump samples collected <0.5 m from dock-hung scallops during the spawning season, and (c) gonad index (GSI, gonad weight/soft tissue weight) from a nearby population during spawning season. Error bars are ± 1 SD. Full moons are indicated by open circles. Only gene copy numbers are reported in (b) due to the unknown cell concentration of the samples

that have been reported for these 2 species in GenBank (e.g. Wang et al. 2006). Overall, our phylogenetic reconstruction establishes the homology of the ITS sequences we generated with publicly available molluscan sequences. It suggests that the ITS region may serve as an informative locus for the development of eDNA assays for several species of commercially important bivalves in Maine coastal waters.




Fig. 4. Bayesian phylogram with proportional branch lengths inferred from ITS sequences of scallops, mussels and oysters. Full length ITS sequences from this study are highlighted in blue and include GenBank accession number plus clone ID. The scale bar represents 0.4 substitutions per site in the underlying DNA sequence alignment

4. DISCUSSION

Environmental DNA samples from aquatic ecosystems can contain genetic material from tens of thousands of species, and have the potential to inform ecological studies across multiple trophic levels. As eDNA science rapidly emerges as an area of interdisciplinary research, it is proving useful in identifying and quantifying a spectrum of biota, from microbes (e.g. Countway & Caron 2006) to metazoans, including life-stages from larvae to adults (e.g. Fox et al. 2005, Thomsen et al. 2012, Pikitch 2018). These data can be used to infer reproductive events through aggregation activity (Tillotson et al.

2018), behavior (Erickson et al. 2016, de Souza et al. 2016), and community structure (Rees et al. 2014).

Size-fractionation has been used to detect an eDNA signal (via qPCR) for the extremely small (~0.8 μm) microbial eukaryote, *Ostreococcus*, in 2 different size-classes of plankton (i.e. 0.7–80 μm vs. 80–200 μm), revealing the overwhelming presence of *Ostreococcus* in the smaller size-class (Countway & Caron 2006). In a situation analogous to the present study, the work by Countway & Caron (2006) demonstrated a dramatic increase in gene copy numbers with the smaller size-class of plankton, indicating a large bloom of *Ostreococcus*. Size-fractionation of plankton samples has proved to be a useful technique to explore functional differences among marine organisms in certain size-classes. To our knowledge, this is the first time size-fractionation combined with a quantitative eDNA assay has been applied to the detection of broadcast spawning events in the marine environment.

Our DNA dilution series of ITS clones indicates a reliable relationship between Cq values from scallop sperm cell extracts and ITS copy numbers using the TaqMan probe and primer set that we developed for *Placopecten magellanicus* (Table 1). The application of the dilution series to our scallop sperm samples suggests that not all of the scallop sperm cells were lysed during the DNA extraction process, or conversely the cells may have been under-

counted by our hemocytometer estimates. Another possibility is that our TaqMan probe and primers do not bind as well to potential (but unknown) variations of the *P. magellanicus* ITS region. It is likely that there are other variants of the ITS region in the *P. magellanicus* genome, given that all 8 of the sequences in our study were unique. In fact, the genome of *P. magellanicus* is thought to be among the largest of the scallop species (Krause & von Brand 2016, and references therein). It is generally accepted that species with larger genomes have more copies of the rRNA operon (Prokopowich et al. 2003) with potentially greater numbers of ITS sequence variants. We attempted to control for this unknown

sequence variability by designing the scallop qPCR assay in a region that was conserved for all of the ITS sequences in our study. Specifically, the 211 bp region that we targeted for qPCR amplification showed very little DNA sequence variation among the 8 *P. magellanicus* ITS clones and the single *P. magellanicus* transcriptome sequence (Pairett & Serb 2013) from GenBank. Notably, the GenBank sequence displayed 100% homology to the consensus sequence of our 8 ITS clones within the qPCR region. Only 3 single-nucleotide variants were observed among 4 of the *P. magellanicus* ITS sequences between the forward and reverse qPCR primers, and none of these variations were found within the probe or primer regions. Our work to characterize the *P. magellanicus* ITS region via low-throughput cloning and sequencing has generated the first full-length ITS sequences for this species. Although these sequences likely do not capture all of the diversity in the ITS region of *P. magellanicus*, they were extremely useful for development of our novel quantitative eDNA application. It is probable that high-throughput DNA sequencing could be used to address the question of ITS sequence variation, given the recent advances in long-read, next generation sequencing technology for DNA barcoding (Hebert et al. 2018).

The GoTaq[®] G2 Hot Start DNA polymerase that was used to amplify the ITS region has an error rate that is consistent with values reported previously for Taq DNA polymerase (Promega technical support pers. comm.) ranging from 20×10^{-6} to 100×10^{-6} errors per nucleotide incorporated (Sambrook & Russell 2001). Although it is likely that some of the variable positions in the ITS region were the result of PCR error, a large proportion of the variants were repeated in multiple sequences (Fig. S1), suggesting that much of the variability across the full-length ITS region (~750 bp) reflected true genetic differences. Errors in the ITS sequences caused by PCR might have contributed to slight variations in the topology of the phylogenetic tree, but they were unlikely to have impacted the qPCR assay due to the high degree of sequence similarity that was observed in the targeted region for the 8 clone sequences and 1 transcriptome sequence. Despite the observed offset between cell counts and gene copy numbers, the strong linear relationship (Fig. 2) suggests that any possible errors related to cell lysis, cell abundance estimates, or undetected variation in the ITS regions were consistent across all samples. It is possible that some sea scallop sperm cells released during spawns are anucleated, as observed in the protan-

dric hermaphroditic bivalves *Tellimya ferruginosa*, *T. tenella*, and *Montacuta percompressa* (Fox et al. 2007). However, anucleated paraspermatazoa have not been observed in scallops to the best of our knowledge.

The results from our dockside experiment suggest that we detected at least 2 male spawning events in the field, both during periods of time when decreases in GSI occurred (Fig. 4). While particles >20 μm showed a spike in the abundance of scallop eDNA at the same time as the first spike in scallop eDNA for particles <20 μm , this parallel response did not happen during the second spike in scallop eDNA for particles <20 μm . We cannot confirm whether the larger particles were eggs or were attributed to some other source of scallop eDNA. It is possible that scallop eDNA in the larger size-fraction could have originated from other scallop tissue due to decaying adult biomass. However, we did not observe any scallop mortality during the dockside experiment, so it is unlikely that cellular debris could have originated from such an event. Cell buoyancy could also impact our eDNA signal, causing gametes to sink to greater depths or float to the surface. While sperm are usually neutrally buoyant, they have been found to swim downwards across multiple phyla (Falkenberg et al. 2016) which may impact eDNA signals. Additionally, sea scallop eggs are negatively buoyant (S. R. Bayer & R. A. Wahle unpubl. data) but have been observed to travel >3 m in unidirectional flumes without sinking (S. R. Bayer pers. obs.). This last observation suggests that the interaction between gametes and the surrounding physical dynamics, including flow rates, turbulence, and shear (i.e. Denny & Shibata 1989, Riffell & Zimmer 2007), deserves greater study and attention when evaluating eDNA data.

A factor to consider in our observations is persistence of the spawning signal. Within the Damariscotta River tidal estuary, we predict that an eDNA signal from spawning would be impacted the most during maximum ebb and flood tides, showing up many kilometers downstream from the adult source, similar to the results of the O'Donnell et al. (2017) study. During these periods of high flow, we would expect to have challenges detecting accurate spatiotemporal eDNA signals similar to that of river and stream studies (i.e. Jerde et al. 2016, Shogren et al. 2017). We sampled specifically at slack tides to avoid these high-flow conditions that may sweep away scallop gametes rapidly. This ensured, to the best of our ability, observation of a signal that was as physically close to our source population as possible.

Both of our observed spikes in sperm coincided with the first quarter of the moon (see Fig. 3). Lunar cues have been documented in sea scallops (Bonardelli et al. 1996). In the Damariscotta River, we have consistently observed female scallop spawning on the third quarter of the moon (S. R. Bayer pers. obs.). Our preliminary data suggest that the temporal spawning patterns in males and females may be worthy of closer investigation given the possible mismatch of male and female spawning events observed in our time-series. Given that lunar cycles strongly influence tides, tidally induced variability in flow may be an important spawning trigger for sea scallops, but the proximate mechanism is not understood. High-flow environments can reduce gamete concentration and wash a majority of sperm and eggs away from spawning adults, while low-flow conditions can allow for higher fertilization success (e.g. Denny & Shibata 1989, Coma & Lasker 1997, Marshall 2002, Riffell & Zimmer 2007).

We do not know how quickly *P. magellanicus* gametes degrade in natural marine conditions. However, we do know that sea scallop eggs remain viable for fertilization between 8 and 24 h, while sperm may be viable for several hours (Bayer et al. 2016). It is assumed sperm DNA persists for several hours while a sperm swims and searches for an egg to fertilize, but this may vary with acidity, salinity and temperature of their environment (Desrosiers et al. 1996). Very few studies investigating eDNA dynamics have accounted for eDNA degradation and the persistence of microbial eDNA ranges from 1 d to >2 wk in aquatic environments (Barnes et al. 2014). Degradation and persistence should be given serious consideration when using eDNA data in fishery and conservation management decisions.

Accurate identification of invertebrate gametes and larvae can be extraordinarily difficult if not impossible by microscopy. Creating a species-specific qPCR assay to detect scallop eDNA allowed us to monitor and quantify sea scallop spawning events and could eventually lead to estimates of the size of subsequent larval output from water samples taken *in situ*. Application of our probe and primers to plankton samples collected near scallop beds would help indicate the presence, magnitude, and timing of scallop spawning events. Collection of size-fractionated samples would be a necessary step to help distinguish sperm cells from eggs, embryos and larvae (~10-fold the size of sperm; Naidu 1970). Expanding beyond scallops, this gamete detection assay could help identify the presence of rarer species through spawning events. Collected eDNA has been used to

identify the presence of rare or elusive species in aquatic environments, like sturgeon Acipenseridae (Pfleger et al. 2016). However, Erickson et al. (2016) found no relationship between eDNA and spawning activity of bigheaded carps (*Hypophthalmichthys* spp.) in rivers. This is further evidence that hydrography is to be carefully considered and studied when studying spawning events.

The role of sperm limitation in population dynamics of free-spawning species has been a long-standing question in marine ecology (Petersen & Levitan 2001), particularly for species whose populations have been commercially exploited or experienced high levels of natural mortality (Yund 2000, Petersen & Levitan 2001, Gascoigne & Lipcius 2004). The challenge has been to detect and quantify gametes in wild populations as they are spawned. One approach has been to use time-integrated fertilization assays (Levitan et al. 1992, Levitan & Young 1995, Wahle & Peckham 1999, Gaudette et al. 2006, Bayer et al. 2016, 2018), a labor- and time-intensive method that requires the induced spawning in the laboratory of viable unfertilized eggs prior to their deployment *in situ* within artificial mesh containers that are permeable to sperm. The method is tractable for a small number of species, but is prone to experimental artifacts (e.g. Levitan 1991, Levitan et al. 1992, Styan 1998a, Bayer et al. 2016, 2018). Development of a molecular assay could dramatically enhance the measurement of spawning events and aid our understanding of spawning and fertilization dynamics in nature.

Our probe and primers are species-specific for *P. magellanicus*, and could aid in quantifying other water-borne life stages in addition to sperm, such as eggs, embryos and larvae. Our field experiment demonstrated that the scallop qPCR assay was able to detect sperm *in situ*. In future studies, we would like to rigorously develop methods for reliably detecting and quantifying eggs and embryos, accounting for physical properties like degradation and buoyancy. Another way to augment this kind of study would be to determine relative mitochondrial eDNA and nuclear eDNA abundances of sea scallops to detect gametes. This method has been demonstrated as reliable in the endangered Macquarie perch *Macquaria australasica* (Bylemans et al. 2017) but not necessarily invertebrates to the best of our knowledge. Identifying gamete and embryo presence and abundance has influenced stock assessments for cod (*Gadus morhua*) in the Irish Sea using similar molecular methods (Fox et al. 2008). Several larval parasitic copepod species, which are difficult to differentiate

by eye, were accurately identified and quantified in plankton samples by a TaqMan probe and primer set (McBeath et al. 2006). Similarly, larval stages of bivalve species can appear almost identical in appearance during early development (Garland & Zimmer 2002). Given that gametes and larvae of marine organisms can be difficult to distinguish, qPCR analysis could revolutionize the identification and quantification of species-specific spawning events in the field. This analysis may also determine if larval cohorts are found within the boundaries of critical fishing grounds and could greatly influence fishery management decisions and the designation of area closures. This method has the potential to be far more practical and less artifact-prone than the use of time-integrated fertilization assays. It could also improve detection of spawning events in remote and hard to access locations that do not lend themselves to fertilization assays, such as offshore and deep-water populations. To build on the specific scallop qPCR assay we developed with this study, future investigations with this novel eDNA methodology should involve the development of probes and primers for other broadcast spawning invertebrates and involve more rigorous field tests around naturally spawning populations.

Acknowledgements. The research was financially supported by 2 Maine Sea Grant program development grants DV-16-06 and DV-17-12 awarded to S.R.B., P.D.C. and R.A.W., a University of Maine Janet Waldron Doctoral Research Fellowship, NSF grant OCE1333797 that supported S.R.B. while working in the Yund Lab, and support from Bigelow Laboratory for Ocean Sciences to P.C. We thank the Marine Culture Laboratory at the Darling Marine Center for the use of spawning tables and the filter room. We also thank S. Pearson for aiding S.R.B. in filtering technique, and the Rich laboratory for allowing us to temporarily use their space and deep freezer at the Darling Marine Center. We thank NSF REU intern Catherine Carlisle and lab technician Nick Marquis for their help in DNA extractions in the Countway laboratory. Scallops were collected by divers R.A.W., B. Favitta, and C. Huntsberger. Dock sampling was made possible by the plankton pump provided by P. Yund. S. Peart, K. Stamieszkin and T. Young-Bayer all helped with dockside sampling. Bigelow Laboratory's Center for Venture Research - Seafood Solutions, helped to offset publication costs.

LITERATURE CITED

- Ahlgren NA, Rocap G (2012) Diversity and distribution of marine *Synechococcus*: Multiple gene phylogenies for consensus classification and development of qPCR assays for sensitive measurement of clades in the ocean. *Front Microbiol* 3:213
- Allee WC (1931) Animal aggregations. A study in general sociology. University of Chicago Press, Chicago, IL
- Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25:3389–3402
- Andersen P, Thronsdon J (2004) Estimating cell numbers. In: Hallegraeff GM, Anderson DM, Cembella AD (eds) Manual on harmful marine microalgae. UNESCO, Paris, p 99–129
- Andree KB, Fernández-Tejedor M, Elandaloussi LM, Quijano-Scheggia S and others (2011) Quantitative PCR coupled with melt curve analysis for detection of selected *Pseudo-nitzschia* spp. (Bacillariophyceae) from the northwestern Mediterranean Sea. *Appl Environ Microbiol* 77:1651–1659
- Babcock RC, Mundy CN (1992) Reproductive biology, spawning and field fertilization rates of *Acanthaster planci*. *Mar Freshw Res* 43:525–533
- Babcock R, Mundy C, Keesing J, Oliver J (1992) Predictable and unpredictable spawning events: in situ behavioural data from free-spawning coral reef invertebrates. *Invertebr Reprod Dev* 22:213–227
- Barnes MA, Turner CR, Jerde CL, Renshaw MA, Chadderton WL, Lodge DM (2014) Environmental conditions influence eDNA persistence in aquatic systems. *Environ Sci Technol* 48:1819–1827
- Bayer SR, Wahle RA, Jumars PA, Brady DC (2016) Measuring scallop fertilization success in the field: chamber design and tests. *Mar Ecol Prog Ser* 551:141–154
- Bayer SR, Wahle RA, Brady DC, Stokesbury KDE, Jumars PA, Carey J (2018) Fertilization dynamics in scallop aggregations: reconciling model predictions and field measurements of density effects. *Ecosphere* 9:e02359
- Benson DA, Cavanaugh M, Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J, Sayers EW (2013) GenBank. *Nucleic Acids Res* 41:36–42
- Bonardelli JC, Himmelman JH, Drinkwater K (1996) Relation of spawning of the giant scallop, *Placopecten magellanicus*, to temperature fluctuations during downwelling events. *Mar Biol* 124:637–649
- Brazeau DA, Lasker HR (1992) Reproductive success in the Caribbean octocoral *Briareum asbestinum*. *Mar Biol* 114:157–163
- Bylemans J, Furlan EM, Hardy CM, McGuffie P, Lintermans M, Gleeson DM (2017) An environmental DNA-based method for monitoring spawning activity: a case study, using the endangered Macquarie perch (*Macquaria australasica*). *Methods Ecol Evol* 8:646–655
- Carey JD, Wahle RA, Stokesbury KDE (2013) Spatial scaling of juvenile–adult associations in northwest Atlantic sea scallop *Placopecten magellanicus* populations. *Mar Ecol Prog Ser* 493:185–194
- Coma R, Lasker HR (1997) Effects of spatial distribution and reproductive biology on in situ fertilization rates of a broadcast-spawning invertebrate. *Biol Bull* 193:20–29
- Countway PD, Caron DA (2006) Abundance and distribution of *Ostreococcus* sp. in the San Pedro Channel, California, as revealed by quantitative PCR. *Appl Environ Microbiol* 72:2496–2506
- de Souza LS, Godwin JC, Renshaw MA, Larson E (2016) Environmental DNA (eDNA) detection probability is influenced by seasonal activity of organisms. *PLOS ONE* 11:e0165273
- Denny M, Shibata MF (1989) Consequences of surf-zone turbulence for settlement and external fertilization. *Am Nat* 134:859–889
- Desrosiers RR, Désilets J, Dubé F (1996) Early developmental events following fertilization in the giant scallop *Pla-*

- copecten magellanicus*. Can J Fish Aquat Sci 53: 1382–1392
- ✦ Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res 32:1792–1797
- ✦ Erickson RA, Rees CB, Coulter AA, Merkes CM and others (2016) Detecting the movement and spawning activity of bigheaded carps with environmental DNA. Mol Ecol Resour 16:957–965
- ✦ Falkenberg, LJ, Havenhand, JN, Styan, CA (2016) Sperm accumulated against surface: a novel alternative bioassay for environmental monitoring. Mar Environ Res 114: 51–57
- ✦ Forootan A, Sjöback R, Björkman J, Sjögreen B, Linz L, Kubista M (2017) Methods to determine limit of detection and limit of quantification in quantitative real-time PCR (qPCR). Biomol Detect Quantif 12:1–6
- ✦ Fox CJ, Taylor MI, Pereyra R, Villasana MI, Rico C (2005) TaqMan DNA technology confirms likely overestimation of cod (*Gadus morhua* L.) egg abundance in the Irish Sea: implications for the assessment of the cod stock and mapping of spawning areas using egg based methods. Mol Ecol 14:879–884
- ✦ Fox TH, Jespersen Å, Lützen J (2007) Sperm transfer and reproductive biology in species of hermaphroditic bivalves (Galeommatoida: Montacutidae). J Morphol 268: 936–952
- ✦ Fox CJ, Taylor M, Dickey-Collas M, Fossum P and others (2008) Mapping the spawning grounds of North Sea cod (*Gadus morhua*) by direct and indirect means. Proc R Soc B 275:1543–1548
- ✦ Garland ED, Zimmer CA (2002) Techniques for the identification of bivalve larvae. Mar Ecol Prog Ser 225:299–310
- ✦ Garneau MÉ, Schnetzer A, Countway PD, Jones AC, Seubert EL, Caron DA (2011) Examination of the seasonal dynamics of the toxic dinoflagellate *Alexandrium catenella* at Redondo Beach, California, by quantitative PCR. Appl Environ Microbiol 77:7669–7680
- ✦ Gascoigne J, Lipcius RN (2004) Allee effects in marine systems. Mar Ecol Prog Ser 269:49–59
- ✦ Gaudette J, Wahle RA, Himmelman JH (2006) Spawning events in small and large populations of the green sea urchin *Strongylocentrotus droebachiensis* as recorded using fertilization assays. Limnol Oceanogr 51:1485–1496
- ✦ Hart DR, Rago PJ (2006) Long-term dynamics of U.S. Atlantic sea scallop *Placopecten magellanicus* populations. N Am J Fish Manage 26:490–501
- ✦ Hebert PDN, Braukmann TWA, Prosser SWJ, Ratnasingham S and others (2018) A sequel to Sanger: amplicon sequencing that scales. BMC Genomics 19:219
- ✦ Heid CA, Stevens J, Livak KJ, Williams PM (1996) Real time quantitative PCR. Genome Res 6:986–994
- ✦ Hewson I, Brown JM, Gitlin SA, Doud DF (2011) Nucleopolyhedrovirus detection and distribution in terrestrial, freshwater, and marine habitats of Appledore Island, Gulf of Maine. Microb Ecol 62:48–57
- ✦ Holland PM, Abramson RD, Watson R, Gelfand DH (1991) Detection of specific polymerase chain reaction product by utilizing the 5'→3' exonuclease activity of *Thermus aquaticus* DNA polymerase. Proc Natl Acad Sci USA 88: 7276–7280
- ✦ Huelsenbeck JP, Ronquist F (2001) MRBAYES: Bayesian inference of phylogenetic trees. Bioinformatics 17:754–755
- ✦ Insua A, López-Piñón MJ, Freire R, Méndez J (2003) Sequence analysis of the ribosomal DNA internal transcribed spacer region in some scallop species (Mollusca: Bivalvia: Pectinidae). Genome 46:595–604
- ✦ Jerde CL, Olds BP, Shogren AJ, Andruszkiewicz EA, Mahon AR, Bolster D, Tank JL (2016) Influence of stream bottom substrate on retention and transport of vertebrate environmental DNA. Environ Sci Technol 50:8770–8779
- Krause MK, von Brand E (2016) Chapter 9: scallop genetics and genomes. In: Shumway SE, Parsons GJ (eds) Scallops: biology, ecology, aquaculture, and fisheries, 3rd Edn. Elsevier, Amsterdam
- ✦ Langton RW, Robinson WE, Schick D (1987) Fecundity and reproductive effort of sea scallops *Placopecten magellanicus* from the Gulf of Maine. Mar Ecol Prog Ser 37:19–25
- ✦ Lazoski C, Gusmão J, Boudry P, Solé-Cava AM (2011) Phylogeny and phylogeography of Atlantic oyster species: evolutionary history, limited genetic connectivity and isolation by distance. Mar Ecol Prog Ser 426:197–212
- ✦ Levitan DR (1991) Influence of body size and population density on fertilization success and reproductive output in a free-spawning invertebrate. Biol Bull 181:261–268
- ✦ Levitan DR (1995) The ecology of fertilization in free spawning invertebrates. In: McEdward L (ed) Ecology of marine invertebrate larvae. CRC Press, Boca Raton, FL, p 123–156
- Levitan DR, Petersen C (1995) Sperm limitation in the sea. Trends Ecol Evol 10:228–231
- Levitan DR, Sewell MA (1998) Fertilization success in free-spawning marine invertebrates: review of the evidence and fisheries implications. Can Spec Pub Fish Aquat Sci 125:159–164
- ✦ Levitan DR, Young CM (1995) Reproductive success in large populations: empirical measures and theoretical predictions of fertilization in the sea biscuit *Clypeaster rosaceus*. J Exp Mar Biol Ecol 190:221–241
- ✦ Levitan DR, Sewell MA, Chia FS (1992) How distribution and abundance influence fertilization success in the sea urchin *Strongylocentrotus franciscanus*. Ecology 73: 248–254
- ✦ Marquis ND, Record NR, Robledo JAF (2015) Survey for protozoan parasites in Eastern oysters (*Crassostrea virginica*) from the Gulf of Maine using PCR-based assays. Parasitol Int 64:299–302
- ✦ Marshall DJ (2002) In situ measures of spawning synchrony and fertilization success in an intertidal, free-spawning invertebrate. Mar Ecol Prog Ser 236:113–119
- ✦ Matteson AR, Rowe JM, Ponsero AJ, Pimentel TM, Boyd PW, Wilhelm SW (2013) High abundances of cyanomyoviruses in marine ecosystems demonstrate ecological relevance. FEMS Microbiol Ecol 84:223–234
- ✦ McBeath AJ, Penston MJ, Snow M, Cook PF, Bricknell IR, Cunningham CO (2006) Development and application of real-time PCR for specific detection of *Lepeophtheirus salmonis* and *Caligus elongatus* larvae in Scottish plankton samples. Dis Aquat Org 73:141–150
- ✦ Medlin LK, Elwood HJ, Stickel S, Sogin ML (1988) The characterization of enzymatically amplified eukaryotic 16S-like rRNA-coding regions. Gene 71:491–499
- ✦ Mendo T, Moltschanivskyj N, Lyle JM, Tracey SR, Semmens JM (2014) Role of density in aggregation patterns and synchronization of spawning in the hermaphroditic scallop *Pecten fumatus*. Mar Biol 161:2857–2868
- ✦ Moorthi SD, Countway PD, Stauffer BA, Caron DA (2006) Use of quantitative real-time PCR to investigate the dynamics of the red tide dinoflagellate *Lingulodinium polyedrum*. Microb Ecol 52:136–150

- Myers RA, Barrowman NJ, Hutchings JA, Rosenberg AA (1995) Population dynamics of exploited fish stocks at low population levels. *Science* 269:1106–1108
- Naidu KS (1970) Reproduction and breeding cycle of the giant scallop *Placopecten magellanicus* (Gmelin) in Port au Port Bay, Newfoundland. *Can J Zool* 48:1003–1012
- Nylander JAA (2004) MrModeltest v2. Program distributed by the author. Evolutionary Biology Centre, Uppsala University https://www.researchgate.net/publication/285805344_MrModeltest_V2_Program_Distributed_by_the_Author
- O'Donnell JL, Kelly RP, Shelton AO, Samhuri JF, Lowell NC, Williams GD (2017) Spatial distribution of environmental DNA in a nearshore marine habitat. *PeerJ* 5:e3044
- Oliver J, Babcock R (1992) Aspects of the fertilization ecology of broadcast spawning corals: sperm dilution effects and in situ measurements of fertilization. *Biol Bull* 183:409–417
- Oliverio M, Mariottini P (2001) Contrasting morphological and molecular variation in *Coralliophila meyendorffii* (Muricidae, Coralliophilinae). *J Molluscan Stud* 67:243–246
- Op De Beeck M, Lievens B, Busschaert P, Declerck S, Vangronsveld J, Colpaert JV (2014) Comparison and validation of some ITS primer pairs useful for fungal metabarcoding studies. *PLOS ONE* 9:e97629
- Pairett AN, Serb JM (2013) De novo assembly and characterization of two transcriptomes reveal multiple light-mediated functions in the scallop eye (Bivalvia: Pectinidae). *PLOS ONE* 8:e69852
- Pan M, McBeath AJ, Hay SJ, Pierce GJ, Cunningham CO (2008) Real-time PCR assay for detection and relative quantification of *Liocarcinus depurator* larvae from plankton samples. *Mar Biol* 153:859–870
- Petersen CW, Levitan DR (2001) The Allee effect: a barrier to recovery by exploited species. In: Reynolds JD, Robinson JG (eds) *Conservation of exploited species*. Cambridge University Press, Cambridge, p 281–300
- Petersen CW, Warner RR, Cohen S, Hess HC, Sewell AT (1992) Variable pelagic fertilization success: implications for mate choice and spatial patterns of mating. *Ecology* 73:391–401
- Petersen CW, Warner RR, Shapiro DY, Marconato A (2001) Components of fertilization success in the bluehead wrasse, *Thalassoma bifasciatum*. *Behav Ecol* 12:237–245
- Pfleger MO, Rider SJ, Johnston CE, Janosik AM (2016) Saving the doomed: using eDNA to aid in detection of rare sturgeon for conservation (Acipenseridae). *Glob Ecol Conserv* 8:99–107
- Pikitch EK (2018) A tool for finding rare marine species. *Science* 360:1180–1182
- Prokopowich CD, Gregory TR, Crease TJ (2003) The correlation between rDNA copy number and genome size in eukaryotes. *Genome* 46:48–50
- Reece KS, Dungan CF, Burrenson EM (2008) Molecular epizootiology of *Perkinsus marinus* and *P. chesapeaki* infections among wild oysters and clams in Chesapeake Bay, USA. *Dis Aquat Org* 82:237–248
- Rees HC, Maddison BC, Middleditch DJ, Patmore JRM, Gough KC (2014) The detection of aquatic animal species using environmental DNA—a review of eDNA as a survey tool in ecology. *J Appl Ecol* 51:1450–1459
- Riffell JA, Zimmer RK (2007) Sex and flow: the consequences of fluid shear for sperm–egg interactions. *J Exp Biol* 210:3644–3660
- Rowe S, Hutchings JA, Bekkevold D, Rakitin A (2004) Depensation, probability of fertilization, and the mating system of Atlantic cod (*Gadus morhua* L.). *ICES J Mar Sci* 61:1144–1150
- Salvi D, Macali A, Mariottini P (2014) Molecular phylogenetics and systematics of the bivalve family Ostreidae based on rRNA sequence-structure models and multilocus species tree. *PLOS ONE* 9:e108696
- Sambrook J, Russell DR (2001) *Molecular cloning: a laboratory manual*, 3rd edn. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY
- Schwalbach MS, Fuhrman JA (2005) Wide-ranging abundances of aerobic anoxygenic phototrophic bacteria in the world ocean revealed by epifluorescence microscopy and quantitative PCR. *Limnol Oceanogr* 50:620–628
- Seamone CB, Boulding EG (2011) Aggregation of the northern abalone *Haliotis kamtschatkana* with respect to sex and spawning condition. *J Shell Res* 30:881–888
- Shogren AJ, Tank JL, Andruszkiewicz E, Olds B, Mahon AR, Jerde CL, Bolster D (2017) Controls on eDNA movement in streams: transport, retention, and resuspension. *Sci Rep* 7:5065
- Smith SJ, Rago P (2004) Biological reference points for sea scallops (*Placopecten magellanicus*): the benefits and costs of being nearly sessile. *Can J Fish Aquat Sci* 61:1338–1354
- Stokesbury KDE (1999) Physical and biological variables influencing the spatial distribution of the giant scallop *Placopecten magellanicus*. *J Shellfish Res* 18:315 (abstract only)
- Stokesbury KDE (2012) Stock definition and recruitment: implications for the US sea scallop (*Placopecten magellanicus*) fishery from 2003 to 2011. *Rev Fish Sci* 20:154–164
- Styan CA (1997) Inexpensive and portable sample for collecting eggs of free-spawning marine invertebrates underwater. *Mar Ecol Prog Ser* 150:293–296
- Styan CA (1998a) The reproductive ecology of the scallop, *Chlamys bifrons* in South Australia. PhD dissertation, University of Adelaide
- Styan CA (1998b) Polyspermy, egg size, and the fertilization kinetics of free-spawning marine invertebrates. *Am Nat* 152:290–297
- Styan CA, Butler AJ (2003) Scallop size does not predict amount or rate of induced sperm release. *Mar Freshwat Behav Physiol* 36:59–65
- Suzuki MT, Taylor LT, DeLong EF (2000) Quantitative analysis of small-subunit rRNA genes in mixed microbial populations via 5'-nuclease assays. *Appl Environ Microbiol* 66:4605–4614
- Swofford DL (2003) PAUP*. Phylogenetic analysis using parsimony (*and other methods). Version 4. Sinauer Associates, Sunderland, MA
- Taylor MI, Fox C, Rico I, Rico C (2002) Species specific TaqMan probes for simultaneous identification of (*Gadus morhua* L.), haddock (*Melanogrammus aeglefinus* L.) and whiting (*Merlangius merlangus* L.). *Mol Ecol Notes* 2:599–601
- Thompson KJ, Inglis SD, Stokesbury KDE (2014) Identifying spawning events of the sea scallop *Placopecten magellanicus* on Georges Bank. *J Shellfish Res* 33:77–87
- Thomsen PF, Kielgast J, Iversen LL, Møller PR, Rasmussen M, Willerslev E (2012) Detection of a diverse marine fish fauna using environmental DNA from seawater samples. *PLOS ONE* 7:e41732

- ✦ Tian RC, Chen C, Stokesbury KDE, Rothschild BJ and others (2009) Dispersal and settlement of sea scallop larvae spawned in the fishery closed areas on Georges Bank. *ICES J Mar Sci* 66:2155–2164
- ✦ Tillotson MD, Kelly RP, Duda JJ, Hoy M, Kralj J, Quinn TP (2018) Concentrations of environmental DNA (eDNA) reflect spawning salmon abundance at fine spatial and temporal scales. *Biol Conserv* 220:1–11
- ✦ Tremblay MJ, Loder JW, Werner FE, Naimie CE, Page FH, Sinclair MM (1994) Drift of sea scallop larvae *Placopecten magellanicus* on Georges Bank: a model study of the roles of mean advection, larval behavior and larval origin. *Deep Sea Res II* 41:7–49
- ✦ Vadopalas B, Bouma JV, Jackels CR, Friedman CS (2006) Application of real-time PCR for simultaneous identification and quantification of larval abalone. *J Exp Mar Biol Ecol* 334:219–228
- ✦ Van der Auwera G, Chapelle S, De Wächter R (1994) Structure of the large ribosomal subunit RNA of *Phytophthora megasperma*, and phylogeny of the oomycetes. *FEBS Lett* 338:133–136
- ✦ Vandersea MW, Kibler SR, Van Sant SB, Tester PA and others (2017) qPCR assays for *Alexandrium fundyense* and *A. ostenfeldii* (Dinophyceae) identified from Alaskan waters and a review of species-specific *Alexandrium* molecular assays. *Phycologia* 56:303–320
- ✦ Vierna J, Martínez-Lage A, González-Tizón AM (2010) Analysis of ITS1 and ITS2 sequences in *Ensis* razor shells: suitability as molecular markers at the population and species levels, and evolution of these ribosomal DNA spacers. *Genome* 53:23–34
- ✦ Vogel H, Czihak G, Chang P, Wolf W (1982) Fertilization kinetics of sea urchin eggs. *Math Biosci* 58:189–216
- ✦ Wahle RA, Peckham SH (1999) Density-related reproductive trade-offs in the green sea urchin, *Strongylocentrotus droebachiensis*. *Mar Biol* 134:127–137
- ✦ Wang S, Bao Z, Zhang L, Li N and others (2006) A new strategy for species identification of planktonic larvae: PCR-RFLP analysis of the internal transcribed spacer region of ribosomal DNA detected by agarose gel electrophoresis or DHPLC. *J Plankton Res* 28:375–384
- ✦ Warner RR, Shapiro DY, Marcanato A, Petersen CW (1995) Sexual conflict: males with highest mating success convey the lowest fertilization benefits to females. *Proc R Soc B* 262:135–139
- ✦ Wong MC, Barbeau MA (2003) Effects of substrate on interactions between juvenile sea scallops (*Placopecten magellanicus* Gmelin) and predatory sea stars (*Asterias vulgaris* Verrill) and rock crabs (*Cancer irroratus* Say). *J Exp Mar Biol Ecol* 287:155–178
- ✦ Yund PO (2000) How severe is sperm limitation in natural populations of marine free-spawners? *Trends Ecol Evol* 15:10–13

Editorial responsibility: Per Palsbøll,
Groningen, The Netherlands

Submitted: October 16, 2018; Accepted: May 10, 2019
Proofs received from author(s): July 3, 2019