

Decadal decline in demersal fish biomass coincident with a prolonged drought and the introduction of an exotic starfish

G. D. Parry*, A. J. Hirst

*Corresponding author: greg.parry@marecol.net.au

Marine Ecology Progress Series 544: 37–52 (2016)




Fig. S1. Annual anomalies in the percentage of hourly wind records at Pt Wilson between 1992–2009: A. $<25 \text{ km.h}^{-1}$, B. $>25 \text{ km.h}^{-1}$, and C. $>40 \text{ km.h}^{-1}$. Dotted lines indicate the difference between the overall mean and means for 1992–96 (pre-drought) and 1997–2010 (drought).




Fig. S2. Annual anomalies in the percentage of east and west components of hourly wind records at Pt Wilson between 1992–2009: A. East component $<25 \text{ km.h}^{-1}$, B. West component $<25 \text{ km.h}^{-1}$, and C. East component $>25 \text{ km.h}^{-1}$, and D. West component $>25 \text{ km.h}^{-1}$. Dotted lines indicate the difference between the overall mean and the mean for 1992–96 (pre-drought) and for 1997–2010 (drought).




Fig. S3. Annual anomalies in the percentage of north and south components of hourly wind records at Pt Wilson between 1992–2009: A. North component <25 km.h⁻¹, B. South component <25 km.h⁻¹, and C. North component >25 km.h⁻¹, and D. South component >25 km.h⁻¹. Dotted lines indicate the difference between the overall mean for 1992–96 (pre-drought) and for 1997–2010 (drought).