

## Species and sexual differences in human-oriented behavior of anemonefish at Okinawa Island, Japan

K. Hayashi\*, K. Tachihara, J. D. Reimer

\*Corresponding author: aetobatu@gmail.com

Marine Ecology Progress Series 616: 219–224 (2019)

Table S1. Number of anemonefish pairs used in the experiments. Site numbers correspond to S1. Species composition differ among sites (Hayashi et al. 2018).

(A) Number of anemonefish pairs used for the experiment 'the effect of approach distance'.

Site No.	Anemonefish species			
	<i>Amphiprion ocellaris</i>	<i>A. clarkii</i>	<i>A. perideraion</i>	<i>A. frenatus</i>
1	2	0	5	0
2	11	3	5	1
3	1	2	5	2
4	1	5	0	6
5	2	5	1	4
6	0	2	0	2
Total	17	17	16	15

(B) Number of anemonefish pairs used for the experiment 'the effect of elapsed time'.

Site No.	Anemonefish species			
	<i>Amphiprion ocellaris</i>	<i>A. clarkii</i>	<i>A. perideraion</i>	<i>A. frenatus</i>
1	2	0	5	0
2	6	3	3	1
3	0	0	0	0
4	0	0	0	0
5	2	3	1	6
6	0	2	0	1
Total	10	8	9	8




Fig. S1. Six study sites set on the western coast of Okinawa-jima Island. Detailed habitat data are shown in Hayashi et al. (2018).